

Effect of organic and inorganic substances on plant growth and survival in khirni (*Manilkara hexandra*)

N. Niranjana and S. R. Barkule

College of Agriculture, Vasant Rao Naik Marathwada Krishi Vidyapeeth, Parbhani, Maharashtra, India

ABSTRACT

The study was carried out to find out the effect of organic and inorganics on growth and survival of khirni (*Manilkara hexandra* L.) at Vasant Rao Naik Marathwada Krishi Vidyapeeth, Parbhani, during 2023-2024. Among treatment combinations, seeds soaked in cow dung slurry for 24 hr showed better stem diameter (3.18 mm), seedling vigour index I (3122.55) and seedling vigour index II (107.92), survival (93.42 %) and lesser incidence of damping off (6.58 %), whereas seedling height (18.04 cm), number of leaves (9.97), leaf area (21.36 cm²), plant fresh weight (3.09 g) and plant dry weight (1.44 g) were observed best in treatment seeds soaked in GA₃ 300 ppm for 12 hr.

Key words: Cow dung, Cow urine, Gibberellic acid, Thiourea, Potassium nitrate.

Khirni (*Manilkara hexandra* L.) belonging to the family Sapotaceae, is a native of tropical South-East Asia. The major khirni growing states in our country are Madhya Pradesh, Gujarat, Rajasthan, Karnataka, Maharashtra and Tamil Nadu. It is mainly propagated by seeds. It is drought hardy and slow growing species commonly grown in laterite soil. It is used as rootstock for sapota propagation.

The non-availability of sufficient rootstock of khirni in southern region of Maharashtra is a limiting factor and it is due to less germination percentage, slow growth of seedlings to attain the graftable size. An attempt has been made to improve the germination of its seeds and subsequent growth. The use of bioregulators in enhancing seed germination and seedling growth of numerous plant species is well known (Malshe *et al.* 2014). Besides, synthetic chemicals and other naturally available bio-products of organics (cow dung and cow urine) are known to contain vital plant growth substance, which enhance the growth and development of plants. Keeping in view, an experiment was conducted to see effect of different chemicals, plant growth regulators, cow-dung slurry and cow urine on germination, growth and development of khirni seeds (Shinde and Malshe, 2015).

The seeds were sown in 20 cm x 15 cm size polythene bags. They were filled with potting mixture, *ie.* soil, FYM and sand (2:1:1 ratio) and uniform sized healthy pre-treated khirni seeds by organics and inorganic substances were sown in bags which were then properly labelled with tags for observation. The Randomized Block Design with 13 treatments and 3 replications was used. The treatments were T₁: seed soaking in cow urine (20 %) for

12 hrs, T₂: seed soaking in cow urine (20 %) for 24 hrs, T₃: seed soaking in cow dung slurry for 12 hrs (1:2 cow dung + water), T₄: seed soaking in cow dung slurry for 24 hrs (1:2 cow dung + water), T₅: seed soaking in cow urine (20 %) for 12 hrs + cow dung slurry for 12 hrs, T₆: seed soaking in GA₃ @ 100 ppm for 12 hrs, T₇: seed soaking in GA₃ @ 200 ppm for 12 hrs, T₈: seed soaking in GA₃ @ 300 ppm for 12 hrs, T₉: seed soaking in KNO₃ (1 %) for 12 hrs, T₁₀: seed soaking in KNO₃ (2 %) for 12 hrs, T₁₁: seed soaking in Thiourea (1 %) 12 hrs, T₁₂: seed soaking in Thiourea (2 %) for 12 hrs, T₁₃: Control (without soaking).

Observations on growth and survival percentage were taken. The statistical analysis was done by following the ANOVA technique (Panse and Sukhatme, 1967). The data pertaining on seedling height of khirni at 60 and 180 days after sowing influenced by different organics and chemicals. There was significant effect of chemicals and organics on seedling height (Table 1). At 60 days after sowing maximum seedling height (6.01 cm) was recorded in treatment T₄ (Seed soaking in cow dung slurry for 24 hrs) which was statistically at par with treatment T₈ (5.56 cm) and T₅ (5.49 cm), while minimum seedling height (4.24 cm) was recorded in treatment T₁₃ (control). At 180 DAS, maximum seedling height (18.04 cm) was recorded in T₈ (seed soaking in GA₃ @ 300 ppm for 12 hrs) which was statistically at par with T₇ (17.57 cm), T₄ (17.50 cm) and T₅ (17.13 cm), while minimum seedling height (11.26 cm) was recorded in treatment T₁₃ (control).

The increase in seedling height was due to effect of GA₃ which stimulates vegetative growth by cell multiplication and cell elongation. It is also reported by Lockhart (1960) that GA₃ induces plasticity in cell wall

*Corresponding author: niranjan93807@gmail.com

Table . 1: Effect of organic and inorganic substances on plant growth and survival in khirni.

Treatment	Seedling height (cm)		Stem diameter (mm)		Number of leaves		Leaf area (cm ²)		Plant fresh weight (g)	Plant dry weight (g)	Seedling Vigour index I	Seedling Vigour index II	Survival (%)	Incidence of damping off (%)
	60 DAS	180 DAS	60 DAS	180 DAS	60 DAS	180 DAS	60 DAS	180 DAS						
	T ₁	5.12	16.4	1.25	2.74	2.33	7.42	2.85						
T ₂	5.20	16.49	1.26	2.78	2.25	7.42	2.92	18.36	2.27	0.98	2597.74	67.24	81.69	18.31
T ₃	5.36	16.82	1.28	2.8	2.51	8.60	3.05	19.75	2.9	1.29	2688.32	90.21	88.04	11.96
T ₄	6.01	17.50	1.43	3.18	2.92	9.22	3.39	20.43	3.01	1.41	3122.55	107.92	93.42	6.58
T ₅	5.49	17.13	1.31	2.85	2.60	8.26	3.13	19.27	2.49	1.10	3046.26	79.00	88.27	11.73
T ₆	4.95	16.14	1.23	2.81	2.11	8.15	2.70	18.83	2.38	1.06	2361.06	65.57	82.17	17.83
T ₇	5.06	17.57	1.24	2.88	2.22	8.49	2.80	19.42	2.74	1.16	2518.24	73.26	85.18	14.82
T ₈	5.56	18.04	1.35	3.02	2.72	9.97	3.21	21.36	3.09	1.44	3073.94	106.28	90.72	9.28
T ₉	4.90	15.45	1.21	2.68	2.01	7.09	2.66	17.5	1.89	0.83	1987.52	45.83	75.71	24.29
T ₁₀	4.93	15.73	1.22	2.71	2.06	7.20	2.68	17.75	2.05	0.88	2195.3	52.76	80.50	19.5
T ₁₁	4.78	14.89	1.19	2.67	1.55	6.42	2.48	16.92	1.68	0.70	1815.41	35.87	73.89	26.11
T ₁₂	4.81	15.14	1.21	2.68	2.00	6.87	2.59	17.14	1.79	0.79	1901.65	42.14	77.89	22.11
T ₁₃	4.24	11.26	1.02	2.31	1.33	6.31	2.28	16.00	1.52	0.58	1434.56	27.88	71.67	28.33
Mean	5.11	16.04	1.24	2.77	2.20	7.80	2.83	18.45	2.3	1.01	2399.74	65.58	82.33	17.67
SEm(±)	0.22	0.37	0.05	0.12	0.14	0.50	0.18	0.66	0.13	0.05	132.09	4.43	3.55	1.33
CD (5%)	0.63	1.08	0.15	0.36	0.41	1.46	0.53	1.93	0.38	0.15	385.57	12.92	10.36	3.88

similar to that of auxins. Thus, it creates water diffusion pressure deficit which results in water-uptake, causing cell elongation and increasing seedling height. These results align with those of Vachhani *et al.* (2014) and Dilip *et al.* (2017).

At 60 DAS, T₄ (seed soaking in cow dung slurry for 24 hrs) recorded maximum number of leaves/plant (2.92) which was statistically at par with T₈ (2.72), T₅ (2.60) and T₃ (2.51). The lowest number of leaves/plant (1.33) was recorded in treatment T₁₃ (control). The treatment T₈ (seed soaking in GA₃ @ 300 ppm for 12 hrs) showed highest number of leaves/plant (9.97) at 180 DAS, which was statistically at par with treatment T₄ (9.92) and T₄ (8.60). The least number of leaves/plant (6.31) was in treatment T₁₃ (control). The maximum number of leaves was in GA₃ treatment due to its presence at apical meristem which results in more synthesis of nucleoprotein responsible for increasing leaf initiation (Sen and Gunthi, 1976). Similar findings were recorded by Sable and Waskar (2009) and Vachhani *et al.* (2014).

At 60 DAS maximum leaf area (3.39 cm²) was recorded in T₄ (seed soaking in cow dung slurry for 24 hrs) which was statistically at par with T₈ (3.21 cm²), T₅ (3.13 cm²), T₃ (3.05 cm²) and T₂ (2.92 cm²). The minimum leaf area (2.28 cm²) was recorded in treatment T₁₃ (control). At 180 days after seed sowing maximum seedling height (21.36 cm²) was recorded in T₈ (seed soaking in GA₃ @ 300 ppm for 12 hrs) which was statistically at par with T₄ (20.43 cm²) and T₃ (19.75 cm²). The minimum leaf area (16.00 cm²) was recorded in treatment T₁₃ (control). The

increase in leaf area of khirni is due to seeds treated with GA₃. The activity of GA₃ at the apical meristem resulting in more synthesis and accumulation of nucleoprotein responsible for increasing leaf initiation and leaf area expansion (Sen and Gunthi, 1976). The similar findings were recorded with Vachhani *et al.* (2014).

The stem diameter (1.43 mm) at 60 days from seed sowing was maximum in T₄ (seed soaking in cow dung slurry for 24 hrs) which was statistically at par with T₈ (1.35 mm), T₅ (1.31 mm) and T₃ (1.28 mm), while minimum stem diameter (1.02 mm) was recorded in T₁₃ (control). Similar trend was observed for stem diameter at 180 days after sowing seed. The increase in stem diameter is due to presence of nitrogen, phosphorus and potassium, essential for plant growth (Khan *et al.* 2010) in cowdung. It was found that increased in stem girth was related to increase in leaf area. It seemed possible that increase in leaf area increased photosynthetic activities and the photosynthates might have been utilized in increasing girth of stem (Mishra and Varma 1980).

The maximum plant fresh weight (3.09 g) at 180 days after sowing was found in T₈ (seed soaking in GA₃ @ 300 ppm for 12 hrs) which was at par with T₄ (3.01 g), T₃ (2.90 g), while minimum plant fresh weight (1.52 g) was recorded under T₁₃ (control).

The increase in fresh weight was observed in seedling treated with GA₃ which resulted in mobilization of water and nutrients transport at higher rate. This might be due to primary cause of stem elongation that enhances more vigour and growth. These results are in confirmation with

those of Bajaniya *et al.* (2018), Sable and Waskar (2009). The maximum plant dry weight was noticed in treatment T₈ (1.44 g) which was at par with T₄ (1.41 g), T₃ (1.29 g), whereas the minimum plant dry weight (0.58 g) was recorded under treatment T₁₃ (control). The maximum seedling vigour index-I (3122.55) at 180 DAS was found in T₄ (seed soaking in cow dung slurry for 24 hrs), which was at par with T₈ (3073.94) and T₅ (3046.26). The minimum seedling vigour index- I (1434.56) was recorded in T₁₃ (control).

At 180 days of sowing maximum seedling vigour index- II (107.92) was found in T₄ (seed soaking in cow dung slurry for 24 hrs), which was at par with treatments T₈ (106.28). The minimum seedling vigour index- II (27.88) was recorded in treatment T₁₃ (control). Seeds treated with cowdung showed the maximum improvement in vigour index of the khirni rootstock. This increase is likely due to the essential nutrients in cow dung such as nitrogen, phosphorus and potassium, essential for plant growth (Khan *et al.* 2010). These nutrients probably enhanced seedling vigour, resulting in better growth characteristics, including more plant height, stem diameter, number of leaves, and leaf area. The similar results were recorded by Desai (1988). The highest survival percentage (93.42 %) was found in T₄ (seed soaking in cow dung slurry for 24 hrs), which was at par with T₈ (90.72 %), T₅ (88.27 %) and T₃ (88.04 %). The least survival percentage (71.67 %) was recorded in T₁₃ (control).

The treatment T₄ (seed soaking in cow dung slurry for 24 hrs) showed lowest incidence of damping off (6.58 %), which was at par with T₈ (9.28 %) seed soaking in GA₃ @ 300 ppm for 12 hrs. The highest incidence of damping off (28.33 %) was recorded in T₁₃ (control).

The increase in survival percentage and decrease of damping percentage is due to seed treated with cowdung which contain nutrients, such as nitrogen, phosphorus and potassium, essential for plant growth (Khan *et al.* 2010) in addition to these nutrients, cow dung contains plant growth-promoting bacteria (PGPB) which can further enhance plant growth (Mukhuba *et al.* 2018). This PGPB live in rhizosphere (area around plant roots) and help promote plant growth by improving nutrient uptake, reducing plant stress and protecting against pathogens (Bashan and De-Bashan, 2005) which increase the vigour of seedlings, which intern increases survival percentage of seedling. The results are in close conformity with findings of Desai (1988) and Patel *et al.* (1996) in khirni.

REFERENCES

- Bajaniya, V. G., Karetha, K. M., Varmore, D. L., Chotaliya, B. M. and Parmar, L. S. (2018). Influence of pre-soaking treatment on seed germination, rooting and survival of khirni (*Manilkara hexandra* L.) seedling cv. Local. *International Journal of Pure & Applied Bioscience* **6**(1): 1629-33.
- Bashan, Y. and De-Bashan, L. E. (2005). Plant growth-promoting. *Encyclopedia of Soils in the Environment* **1**: 103-15.
- Desai, J. D. (1988). 'Seed treatment with cattle-dung and GA and their effect on germination and subsequent growth of seedling of rayan [*Manilkara hexandra* (Roxb)]. M.Sc. (Agric.) Thesis, Gujarat Agricultural University, Anand (unpublished).
- Dilip, W. S., Singh, D., Moharana, D., Rout, S. and Patra, S. S. (2017). Effect of Gibberellic Acid (GA) Different Concentrations at Different Time Intervals on Seed Germination and Seedling Growth of Rangpur Lime. *Journal of Agroecology and Natural Resource Management* **4**(2): 157-65.
- Khan, M.S., Zaidi, A., Ahemad, M., Oves, M. and Wani, P.A. (2010). Plant growth promotion by phosphate solubilizing fungi- current perspective. *Archives of Agronomy and Soil Science* **56**(1): 73-98.
- Lockhart, J. A. (1960). Intracellular mechanism of growth inhibitors radiant energy. *Plant Physiol.* **35**:129.
- Malshe, K. V., Mahadik, S. G., Desai, B. G. and Borate, H. V. (2014). Effect of presowing treatment with gibberellic acid on germination and growth of Khirni (*Manilkara hexandra* L.). *Ann. Plant Physiol.* **28**(2): 73-75.
- Mukhuba, M., Roopnarain, A., Adeleke, R., Moeletsi, M. and Makofane, R. (2018). Comparative assessment of bio-fertiliser quality of cow dung and anaerobic digestion effluent. *Cogent Food and Agriculture* **4**(1): 143-50.
- Sable, P. B. and Waskar, D. P. (2009). Investigation on seed germination and subsequent growth of khirni (*Manilkara hexandra* L.) seedlings. *International J. Tropical Agriculture* **27**(1-2): 33-36.
- Sen, S. K. and Gunthi, P. (1976). Effect of pre-sowing seed treatment on the germination and seedling growth in papaya. *Orissa Journal of Horticulture* **4**: 38- 43.
- Shinde, V. V. and Malshe, K. V. (2015). Effect of cattle urine and cow dung slurry as seed treatment on germination and growth of (*Manilkara hexandra* L.). *J. Ecofriendly Agriculture* **10**(2): 128-30.
- Vachhani, K. B., Gohil, J. H., Pandey, R. and Ray, N. R. (2014). Influence of Chemicals, PGR's and Cow-dung Slurry as Seed Treatment on Germinability, Growth and Development of Khirnee (*Manilkara hexandra* Roxb) under Net House Condition. *Trends in Biosciences* **7**(14): 1641-43.