Current Horticulture 14(1): 31–40, January–March 2026
DOI: 10.5958/2455-7560.2026.00004.X
Received: 02 November 2025; Revised: 5 March 2026; Accepted: 14 March 2026

Next-generation onion (*Allium cepa*) breeding using molecular markers: progress and prospects--a review

Saheb Pal[1*], Labdhi Dedhia[2], Pavithra K.N[1], Narendra Singh[1], Krishna Prakash[1] and Vijay Bahadur Singh Chauhan[1]

[1]*School of Crop Sciences, ICAR-Indian Agricultural Research Institute, Gauria Karma, Barhi, Hazaribagh, Jharkhand, 825 405, India*

Abstract

Onion (*Allium cepa* L.) is an indispensable vegetable crop throughout the globe. It is a member of family Amaryllidaceae (formerly Alliaceae family) and is characterized by 16 somatic chromosomes and a massive 16.4 Gbp genome. Despite its economic significance, the pace of genetic improvement has historically been hindered by the complex and biennial life cycle, photosensitivity, high inbreeding depression and a lack of sufficient genomic resources. The shift from low-throughput markers like RAPD and RFLP to high-resolution and reliable marker systems like SSRs and SNPs has facilitated the construction of saturated linkage maps, leading to identification of QTLs for various traits like resistance to downy mildew, purple blotch, anthracnose and thrips, apart from tolerance to drought stress. Development of specialized mapping populations like doubled haploids has also enabled rapid fixation of heterozygosity and led to identification of responsible loci for different nutritional and bulb-quality traits. Deployment of molecular markers like *PsaO*, *AcPMS1* for rapid identification of Cytoplasmic-Genetic Male Sterility (CGMS) systems, is paving the way for hybrid seed production in a cost-effective manner. Thus, this review synthesizes the evolution of onion breeding, transitioning from conventional phenotypic selection to modern genomic-assisted improvement to overcome the genome complexities for accelerated development of high-yielding, multi-stress-resistant onion cultivars.

Key words: Abiotic stress, *Allium*, Biotic stress, Diseases, Marker-Assisted Selection, Pests, Quality

Downloaded From IP - 216.73.216.217 on dated 02-Nov-2026
Not for Commercial Use
www.indianjournals.com

Onion (*Allium cepa* L.), a member of family Amaryllidaceae (formerly Alliaceae), is characterized by a diploid chromosome complement of 2n = 2x = 16 (Tripathi *et al.,* 2017; Mahala *et al.,* 2024). The primary center of origin is localized to Central Asia-encompassing modern-day Uzbekistan, Turkmenistan, Tajikistan, Afghanistan, Iran and Pakistan (Tripathi and Lawande, 2019). Found growing wild in these regions, *Allium vavilovii* has been identified as the closest wild progenitor to cultivated onion, while Mediterranean basin is recognized as the secondary center of origin.

Globally, onion cultivation is spanned in 5.719 million ha area, yielding a total output of 108.260 million tonnes. Production is heavily concentrated in Asia, which accounts for 64.70% of global volume, followed by Africa (15.70%), Europe (9.70%), and the USA (9.70%). India has emerged as the largest producer of onion followed by China, the USA, Egypt and Bangladesh (FAO STAT, 2024). India produces about 24.266 million tonnes of fresh bulbs from 1.54 million ha area (Anonymous, 2024). It is regarded as the most important export vegetable, contributing to 1.3 million tonnes worth ₹ 2,107.14 crore from India (NHRDF, 2021).

Since the early era of domestication of onion, human being invariably conducted selection leading to the development of different types with respect to bulb shape, size, colour, pungency, photoperiodic response etc. However, despite presence of appreciable variability, speed of onion improvement is not at the pace of other monocot taxa (McCallum, 2007, Varshney *et al.,* 2012). Systematic onion breeding stated with mass selection during the 19th century, followed by discovery of cytoplasmic male sterility (CMS), paving the way for development of F_1 hybrids (Brewster, 2008; McCallum *et al.,* 2008). Presently, F_1 hybrids predominate in long-day season, whereas in short-day conditions, open-pollinated varieties predominate (Brewster, 2008).

The onion breeders following conventional breeding methods often face difficulties because of its high levels of cross-pollination and inbreeding depression, biennial life-cycle, photosensitivity and short seed viability. Genome sequencing poses challenge because of its large 16.4 giga base pairs (Gbp) genome size. Thus, there is an urgent need for deployment of genomic tools to facilitate rapid genetic improvement for yield, quality traits and multi-stress resistance in different genetic backgrounds.

Phenotypic selection of onion often leads to biased results due to the substantial influence of environment on trait expression. Moreover, it is also difficult to improve the traits which have low heritability. Marker-Assisted Selection (MAS) using tightly-linked markers, proves efficient in such cases. The MAS provide additional advantages like high accuracy, rapidity, early trait detection etc. Thus, this review collects the available

Corresponding author : saheb.horti@gmail.com
[2] Division of Flower and Medicinal Crops, ICAR-Indian Institute of Horticultural Research, Hesaraghatta Lake, Bengaluru, Karnataka, 560 089, India

www.indianjournals.com
Not for Commercial Use
Downloaded From IP - 216.73.216.217 on dated 02-Nov-2026

information on progress made by identification and deployment of genomic resources for onion improvement and its prospects.

Genetic Resources

Availability of diverse germplasm is the most crucial factor for a successful breeding programme (Glaszmann *et al.,* 2010). Growing of onion in different agro-ecological zones over centuries led to development of a wide range of landraces, harbouring tremendous genetic variation (Brewster, 2008). However in the late 1980s, development of F_1 hybrids resulted in genetic erosion of those landraces in different countries. However, presently, a huge number of onion accessions are being maintained in different gene banks in the world (Table 1).

Table 1: Major onion gene banks in the world along with their germplasm holdings.

Institute	Location	Number of *Allium* accessions
European Cooperative Programme for Plant Genetic Resources	Germany	14400
Directorate of Onion and Garlic Research	India	2050
N.I. Vavilov Research Institute	Russia	1888
Warwick Crop Center	United Kingdom	1755
National Institute of Agro-biological Sciences	Japan	1352
US Department of Agriculture	USA	1304
Royal Botanic Gardens	United Kingdom	1100
AVRDC – The World Vegetable Center	Taiwan	1082

Despite the wide distribution, the knowledge of genome, population structure and genetic architecture are limited in onion (McCallum, 2007), necessitating genetic resource mining and accelerated genetic analysis (Baldwin *et al.,* 2012a). The genome of onion is very large (~16.4 Gbp) and very few ESTs are available in public domain (McCallum *et al.*, 2001; Kuhl *et al.*, 2004). Forming a partial onion BAC library, Jakše *et al.* (2008) reported genome composition, gene structure and very low gene density of one gene per 168 kb genome. The genomic data and genetic maps of onion and other economically important *Allium* species are available from an online genomic database named as *Allium Map* (https://allium.scinet.org.nz/) (McCallum *et al.*, 2012).

Mapping populations

Predominantly, early generation inbreds are used for development of the mapping populations in onion. The residual heterozygosity in such parental inbreds complicates both marker development and sequence analysis (Baldwin *et al.,* 2012b). In general, F_2 population is most used for molecular mapping owing to difficulties to maintain fertility and plant vigour, while developing Recombinant Inbred Lines (RILs). To overcome this, doubled haploid (DH) onion lines have been developed in both short- and long-day onions (Bohanec, 2009; Baldwin *et al.,* 2012b; Duangjit *et al.,* 2013) for mapping of quantitative and qualitative traits. However, major drawback of doubled haploid accessions is poor seed setting (Bohanec, 2002). In this context, Alan *et al.* (2003 and 2004) developed highly fertile doubled haploid lines from long-day US onion varieties, paving the foundation for better insights into onion genetics and genomics.

Genetic maps/ linkage maps

Linkage Map is a schematic representation of the relative locations of genetic markers present on chromosome of an organism as determined from the frequency of recombination between the marker pairs. The first genetic map of onion consisting of 12 linkage groups was developed by Bradeen and Havey (1995) by using 116 RFLP and RAPD markers in a population developed from BYG15-23 × AC43. To date, this map is regarded as a key reference genetic map of onion. Furthermore, Martin *et al.* (2005) placed 100 additional markers to the same intra-specific map for a better saturation with 14 linkage groups. This map predominantly contained dominant RFLP markers which implies that the large genome size of onion is mainly because of high levels of gene duplication.

Marker systems

RAPD

The RAPD (Randomly Amplified Polymorphic DNA) markers use 10 base long arbitrary primers (decamers) which amplify random segments of genomic DNA in a Polymerase Chain Reaction (PCR). Being random, this marker does not require any sequence information. RAPD had been deployed for characterization of onion germplasm comprising of both wild and cultivated genotypes by Maniruzzaman *et al.* (2010) and Sudha *et al.* (2019).

ISSR

The Inter Simple Sequence Repeat (ISSR) technique amplifies the genomic region falling between two Simple Sequence Repeats (SSRs) by using both anchored and non-anchored primers (Ziętkiewicz *et al.,* 1994). Smolik *et al.* (2007) analyzed genetic diversity of onion genotypes belonging to six different species using 16 ISSR markers. Mukherjee *et al.* (2013) used both RAPD and ISSR for diversity analysis of five *Allium* species and found ISSR to

Table 2: Linkage maps developed in onion along with the marker system utilized

Population type	Population derived from	Marker system employed	Reference(s)
F_2	NW-001 × NW-002	SNP	Jo *et al.* (2017)
F_2	BYG15-23 × AC43	InDel, SNP, SSR, RFLP	Martin *et al.* (2005)
F_2	BYG15-23 × AC43	RFLP, SSR and SNP	King *et al.* (1998); Duangjit *et al.* (2013)
F_2	*Allium cepa* × *Alliumroylei*	AFLP and SSR	Van Heusden *et al.* (2000), McCallum *et al.* (2012)
F_2	W202A × Texas Grano	SSR	McCallum *et al.* (2006, 2007)
F_2	Nasik Red × CUDH2150	SSR and SNP	Baldwin *et al.* (2012b)
F_2	OH1 × 5225	SSR and SNP	Duangjit *et al.* (2013)
F_2	DehyA × B5351C	SNP	Damon and Havey (2014)
F_2	*Allium cepa* × *A. fistulosum*	SSR, InDel, STS and RAPD	Tsukazaki *et al.* (2012)

be more polymorphic between *Allium cepa* and *A. sativum*, while RAPD markers showed better polymorphism within *A. porrum*. In overall, RAPD markers revealed more intraspecific diversity than ISSRs, whereas ISSRs showed more interspecific diversity than RAPD, proving its potential for diversity analysis.

SRAP

The Sequence Related Amplification Polymorphism (SRAP) is a simple marker designed for the amplification of Open Reading Frame (ORF). SRAP primers are generally 17-18 nucleotides long and consist of core sequences (13-14 base pairs) and nucleotides 'CCGG' in the forward primer and 'AATT' in the reverse primer.

RFLP

The Restriction Fragment Length Polymorphism (RFLP) involves a single restriction enzyme which generates fragments of different lengths from the same genomic region of different individuals/strains/lines or species, depending upon the presence or absence of recognition sites. Bark and Havey (1995) deployed this marker system to characterize onion inbred lines.

AFLP

The Amplified Fragment Length Polymorphism (AFLP) was developed by Zabeau and Vos (1993). This technique is a combination of restriction digestion and PCR. Ohara *et al.* (2005) constructed the first genetic linkage map of Japanese bunching onion (*Alliums fistulosum*) using 149 AFLP markers.

SSR

The term *Simple Sequence Repeat* (SSRs) was originally coined by Lilt and Luty (1989) to describe simple sequence fragments generated in PCR. This is also known as Short Tandem Repeat (STR) and Simple Sequence Length Polymorphism (SSLP). The multi-allelic nature and PCR-based detection of the SSR markers makes them more breeder friendly than the SNPs (Pal *et al.,* 2020). More than a hundred SSR markers were developed by Tsukazaki *et al.* (2008) from size-fractionated genomic DNA libraries. SSR makers have provided the greatest insights into onion diversity analysis (Khosa *et al.,* 2014). Mitrová *et al.* (2015) established a panel of 15 easy to score SSR markers which easily differentiated 16 commercial onion cultivars of Czech Republic, making them potential for future diversity analysis studies. Anwar *et al.* (2017) employed 16 SSR and three ISSR markers for diversity analysis in *Allium cepa* L., *A. sativum* L. and *A. kurrat* L. accessions. Ivchenko *et al.* (2017) used six SSRs for identification of onion varieties. While studying the genetic diversity of 16 onion genotypes, using RAPD, SSR and ISSR markers Kesraliker *et al.* (2017), concluded ISSR and SSR to be more reliable over RAPD. Singh *et al.* (2021) and Gupta *et al.* (2022) also deployed microsatellite markers for germplasm characterization and emphasized on these markers over the others for better reproducibility and preciseness.

Single Nucleotide Polymorphism (SNP)

Single Nucleotide Polymorphism **(**SNPs) are one of the most reliable marker systems owing to its preciseness and reproducibility. In recent times, the marker system is gaining importance with the reduction in the sequencing costs. This marker had been deployed for characterization of commercial cultivars by Labate *et al.* (2020), Lee *et al.* (2021) and Jeon *et al.* (2022).

QTL Mapping and Marker Assisted Breeding

Quantitative traits show continuous variation due to their polygenic inheritance and environmental influences. Polygenes produce small individual and cumulative effects on the phenotype. A quantitative trait locus (QTL) is a section of DNA which correlates with the variation of quantitative traits. Thus, QTL mapping employs an algorithm for detection of an association between a phenotypic trait and genetic marker(s).

Disease resistance

Downy mildew, caused by *Peronospora destructor*, is a major yield limiting factor, especially during the *rabi* season. The resistance is conditioned by two dominant genes. Introgression of the resistant genes have been made

www.indianjournals.com
Not for Commercial Use
Downloaded From IP - 216.73.216.217 on dated 02-Nov-2026

Downloaded From IP - 216.73.216.217 on dated 02-Nov-2026
Not for Commercial Use

from *Allium royeli* with the help of tightly-linked AFLP markers are available (Scholten *et al.,* 2007; Khosa *et al.*, 2016). Eidlin *et. al.* (2021) carried out marker-assisted selection using DMR1 marker, linked to the resistant gene '*Pd*', to develop a maintainer line with downy mildew resistance.

Genetics of resistance to *Fusarium* basal rot varies from monogenic to polygenic, depending upon the resistant sources employed (Cramer, 2000, Galvan *et al*., 2008, Taylor *et al.,* 2013). The resistance can also be precisely introduced in susceptible genotypes through CRISPR/Cas mediated genome editing (Pal *et al*., 2025). Incomplete resistance (tolerance) to white rot (Hovius *et al*., 2004) and resistance to pink root (Esfahani and Pour, 2008), *Stemphyllum* blight (Anitha *et al.,* 2011), black mold (Kamal *et al.,* 2012) and Iris Yellow Spot Virus (Bag *et al.,* 2014) has also been identified in onion or other alliums, which are to be deployed in future breeding programmes.

Among biotic stresses, purple blotch causes havoc crop loss ranging from 2.5 to 97% both in bulb and seed crops throughout the globe (Schwartz *et al.,* 2005; Kareem *et al.,* 2012; Tripathi *et al.,* 2013; Nanda *et al.,* 2016; Priya *et al*., 2016). Chand *et al.* (2018) identified *ApR1* governing resistance to this disease in a population derived from Arka Kalyan (resistant parent) and Agrifound Rose (susceptible parent). They further identified one SSR, *i.e*., AcSSR7 and one STS marker, *i.e*., ApR-450 linked to the *ApR1* gene at 1.3 and 1.1 centi Morgan genetic distance, respectively. Recently, KASP based SNP (ApRsnip14, ApRsnip23), SSR (AcSSR7) and SRAP marker (ApR-450) have also been developed for marker-assisted introgression (Sahoo *et al*., 2023) of purple blotch resistance.

Anthracnose disease, caused by *Colletotrichum gloeosporoides*, also poses serious threat to onion cultivation worldwide. Jayaswall *et al.* (2025) shortlisted a total of 131 differentially expressed genes, belonging to gene classes like Mitogen-activated protein kinases, *WRKY* and *MYB* transcription factors, R genes and transcriptional activators, putatively associated with resistance to anthracnose in wild species of onion. Scholten et al. (2016) identified a QTL (*Bs1*) on chromosome 6 for resistance to leaf blight disease caused by *Botrytis squamosa*. Kim *et al.* (2021) identified SCAR-OPAN-1 and SNP-3 HRM markers linked to gray mold disease resistance.

Epicuticular wax for insect resistance

Insects with piercing and sucking type mouth parts recognize their host plants by sensing the organic constituents of epicuticular waxes (Diaz-Montano *et al.,* 2011). Wide natural variation for the amount and types of epicuticular wax has been reported and reduced amount of epicuticular wax (glossy foliage) have been associated with thrips resistance (non-preference) in onion (Bag *et al.,* 2014; Damon and Havey, 2014). Recessive inheritance of reduced epicuticular waxes and found two QTLs each on chromosome 2 and chromosome 5, governing the production of hentriacontanone-16 (primary wax on leaves) and for production of several primary alcohols, respectively (Damon and Havey, 2014). SNP markers are being identified for marker-assisted selection for this trait (Bag *et al.,* 2014; Damon and Havey, 2014).

Drought tolerance

Caldwell *et al.* (2003) investigated soil moisture potential, leaf water potential, growth, photosynthesis, stomatal conductance, leaf transpiration, water use efficiency, membrane function, nature and accumulation of quaternary ammonium compounds in 21 days old onion seedlings which were subjected to various drought regimes by withholding irrigation. They concluded that the onion seedlings were able to tolerate drought by increasing water use efficiency. Though QTL mapping has not been followed in onion for drought tolerance, the related species, *viz. Allium fistulosum* and *A. munzii* and genotypes *viz*., Arka Kalyan, MST 42 and MST 46 have been reported to be drought-tolerant (Singh, 2010), necessitating their deployment in QTL mapping.

Bulb pigmentation

A total of 54 types of flavanoids determine the bulb colour by their relative proportion (Slimestad *et al.,* 2007). However, the major colour compounds are anthocyanin (red), flavonols (pale yellow) and chalcones (bright yellow) (Schwinn *et al.,* 2016). Clarke *et al.* (1944) reported three pairs of genes, *viz.* C, R and I, being involved in the development of pigment in the onion bulbs. *C-c* is a basic color factor and dominant C is necessary to produce any pigment. Consequently, all *cc* plants produce white bulbs. Dominant *R* gene in the presence of *C* produces red pigment and recessive *r* allele is responsible for the production of yellow pigment. The gene, *I* is inhibitory to the coloured bulbs and presence of dominant allele of *I* results in white bulbs. El-Shafie and Davis (1967) reported five genes *viz*., *I, C, G, L, R* (all with two alleles each) responsible for onion bulb colour. Khar *et al.* (2008) developed three different families segregating for bulb colours and identified a SSR marker on chromosome 6 linked to C locus. In cross B2246×B11159, they further reported that red bulbs versus yellow bulbs were controlled by DFR and a locus (*L2*) linked at 6.3 cM to ANS (anthocyanidin synthase). Hence, authors proposed that onions with yellow bulbs were independently selected for numerous times and thus yellow genotypes carry independent mutations in

Downloaded From IP - 216.73.216.217 on dated 02-Nov-2026
Not for Commercial Use
www.indianjournals.com

structural or regulatory genes controlling the production of red bulb colour. Earlier, Kim *et al.* (2004) identified a critical mutation within the chalcone isomerase (CHI) gene that reduces the amount of quercetin, giving rise to gold-coloured bulbs. Further, Kim *et al.* (2005) reported that the inactivation of *DFR* was responsible for colour difference between yellow and red onions and two recessive alleles of *ANS* gene were responsible for the development of pink bulb colour. Later, Park *et al.* (2013) designed functional CAPS marker for differentiating two *DFR-A* alleles *viz.*, DFR-A^{PS} (present in yellow onion) and DFR-A^{DEL} (present in red onion). Tsukazaki *et al.* (2012) developed a linkage map from a population derived from *Allium cepa* and *Allium fistulosum* consisting of 11 linkage groups and total marker coverage of 1040 cM. A QTL for bulb pigmentation was reported on chromosome 7.

Pungency

Bulb pungency is an important determinant of end use of onion bulbs and consumer acceptance. McCallum *et al.* (2007) identified a major QTL for pungency on chromosome 3 by using a $F_{2:3}$ mapping population derived from W202A (pungent type) × Texas Grano 438 (sweet type). Marker assisted transfer of the QTL could be followed for improving pungency into non-pungent genotypes.

Tearless onion

Lachrymatory factor synthase (*LFS*) gene catalyzes the production of the lachrymatory factor in onion bulbs. Masamura *et al.* (2012) mapped a candidate *LFS* gene on chromosome 5 of onion using a complete set of *A. fistulosum*–shallot (*A. cepa* L. *aggregatum* group) monosomic alien addition lines. Eady *et al.* (2008) produced tearless onion by RNAi mediated suppression of *LFS* gene which prevented the conversion of isoalliin into thiosulfinates.

Kato *et al.* (2016) developed non-pungent onions by irradiating seeds with neon-ion at 20 Gy. The level of lachrymatory factor production of the selected bulbs in M_4 generation had 7.5 times lower than those of normal onion, confirmed by the lower mRNA production for the gene coding for Allinase enzyme.

Bulb quality and nutritional traits

Substantial genetic variation for fructo-oligosaccharides, quercetin, epigallocatechin gallate and epicatechin gallate, total and essential amino acids, carbohydrates, total soluble solids, trisulphinates and total antioxidant activity occurs among onion cultivars (Insani *et al.* 2016). The parameters *viz.*, bulb colour, firmness, number of scales, number of growing points, neck thickness, total soluble solids (TSS), pungency and antioxidants are the most important quality determinants in onion among the consumer (Brewster 2008; Goldman 2011. Dry matter, TSS and pungency are positively correlated and show moderate to high heritability, indicating the possibility of simultaneous improvement through selection. However, development of less pungent onion with high dry matter becomes critical (Galamarini *et al.*, 2001; Mallor *et al.,* 2011).

Galmarini *et al.* (2001) used the genetic map of BYG15-23 × AC43 to carry out mapping of loci for pungency, dry matter and TSS. One gene (acid invertase-API89) on chromosome 3 and another gene (phloem unloading sucrose transporter- API66) on chromosome 5, both being responsible for high TSS and dry matter accumulation was reported. Furthermore, the acid invertase gene also elevated pungency, which suggests a possible pleiotropic effect of the gene on bulb composition. Havey *et al.* (2004) conducted further QTL analysis in the same genetic map with extensive analysis of bulb carbohydrates and again reported significant effects of the acid invertase region of chromosome 3 on bulb sucrose content. Three loci each on chromosomes 3,5 and 8 have been identified in a population of *A. cepa* × *A. royeli,* regulating accumulation of non-structural carbohydrates like fructose, fructan, glucose and sucrose (McCallum *et al.,* 2006; Raines *et al.,* 2009), of which *Frc* locus on chromosome 8 is the major determinant for fructan content. This result was later confirmed by Yaguchi *et al.* (2008) by using *Allium fistulosum* and *A. cepa* var. *aggregatum* monosomic alien addition lines. Although two markers *viz.*, ACM033 and ACABE58 have been reported to be linked with the *Frc* locus on chromosome 8, but more tightly linked markers are urgently needed for marker-assisted selection.

Male sterility

Male sterility is the most important genetically controlled pollination control mechanism economizing the hybrid seed production in onion. Cytoplasmic-genetic male sterility (CGMS) system, owing to its ease of maintenance and restoration through the *Ms* locus, is preferred in onion over the CMS. The *Ms* locus codes for penta-trico-peptide repeat (PPR) proteins, leading to restoration of fertility.

The CGMS system has been reported in different cytoplasmic backgrounds of onion like CMS-S (Jones and Emsweller 1936), CMS-T (Berninger 1965) and CMS-T like or CMS-Y cytoplasm (Kim 2014). The fertility restoration of CMS-S cytoplasm gets restored by the fertility restorer, *Ms* locus (Jones and Clarke 1943). However, restoration of fertility in CMS-T cytoplasm is governed by three genes i.e., two complementary genes

Downloaded From IP - 216.73.216.217 on dated 02-Nov-2026
Not for Commercial Use
www.indianjournals.com

and an independent gene (Schweisguth 1973) while in CMS-T like cytoplasm, it is governed by a single nuclear gene (Kim and Kim 2019). Go¨kce *et al.* (2002) and Martin *et al.* (2005), carried out marker-assisted selection using a codominant and PCR-based marker. Manjunathgowda and Selvakumar (2021) validated PsaO marker for its co-dominance and tight linkage to restorer of fertility, *Ms* locus. Thus, the PsaO is suitable rapid identification of desirable recombinants with respect to male fertility while handling segregating generations. Saini et al. (2015) utilized 5'cob and *orfA501* markers, originally reported by Sato et al. (1998) and Engeleke et al. (2003), to distinguish between the male sterile and male fertile cytoplasm of accessions in three different populations. Further, upon testing of two markers *viz.*, OPT and PsaO-linked to *Ms* locus, *OPT* was proven better over the PsaO. Khar et al. (2022) used accD and MKFR markers for *cytotype* identification and AcPMS1 and AcSKP1 markers (Huo *et al.* 1015; Kim *et al.* 2015) to identify *Ms* locus in a diverse germplasm panel. While both accD and MKFR were equally potent for identification of cytoplasm, AcPMS1 was found more reliable to identify the *Ms* locus compared to AcSKP1. Manjunathagowda (2025) utilized two sets of PCR-based molecular markers linked to orf725, originally identified by Kim *et al.* (2009) and reported two short day and Indian accession *viz.* DOGR-MG-2 and DOGR-MG-70 carrying CMS-S cytoplasm.

CONCLUSION AND FUTURE PROSPECTS

Onion is an important vegetable cum spice crop with great impact on the economy across the globe. Though several attempts and significant progress have been made for the genetic improvement, the yield potential is many-a-times affected adversely due to the frequent occurrence of several biotic and abiotic stresses. Moreover, there is differences in the consumer appealing traits for specific end use. Thus, there is an urgent need to develop trait-specific varieties with multiple disease resistance, wider adaptability for varied agro-climatic conditions and suitability for various end use including processing into different products.

Keeping in view the large genome size of onion, there is an urgent need to develop more polymorphic and codominant markers to study genetic variation among the accessions. To meet the increasing demand, development of high-yielding varieties and hybrids in a short span can be achieved by through marker-assisted selection. For this, identification of tightly linked molecular markers and/or identification of candidate genes and functional markers within the gene(s) is urgently needed. Candidate gene association mapping for fructan content and bulb pungency is also required. Mainkar et al. (2023) standardized the genome editing protocol in onion and edited the *AcPDS* gene, paving the way for precise modification of other desirable genes. Keeping in view the dynamic and diverse market segments, less pungent and tearless onion need to be developed through genome editing, apart from development of multi-stress resistant varieties.

DECLARATION

The authors declare that they do not have any conflict of interest.

REFERENCES

Alan, A., Brants, A., Cobb, E., Goldschmied, P., Mutschler, M. and Earle, E. (2004). Fecund gynogenic lines from onion (*Allium cepa* L.) breeding materials. *Plant Sci.* **167**: 1055-66.

Alan, A.R., Mutschler, M.A., Brants, A., Cobb, E and Earle, E.D. (2003). Production of gynogenic plants from hybrids of *Allium cepa* L. and *A. roylei* Stearn. *Plant Sci.* **165**: 1201-11.

Anitha, K.K., Venkateswaran, S.K., Chakrabarty, G., Suresh, Kumar., Babu, S. and Varaprasad, K.S. (2011). Onion genetic resources and pest resistance: the Indian scenario. *Indian J. Plant Prot.* **39**: 81-92.

Anonymous. (2024). Area and Production of Horticultural Crops in India, Final Estimates-2023-2024. Ministry of Agriculture and Farmers' Welfare, Government of India. Accessed from: https://agriwelfare.gov.in/en/StatHortEst; Accessed on: 03/03/2026.

Anwar, G., Helmey, R.K. and Moustafa, M.M. (2017). Genetic diversity among some species of the genus *Allium* L. using SSR and ISSR markers. *Egypt Journal of Genetics and Cytology.* **46**: 221-33.

Bag, S., Schwartz, H.F., Cramer, C.S., Havey, M.J. and Pappu, H.R. (2014). Iris yellow spot virus (Tospovirus: Bunyaviridae): from obscurity to research priority. *Mol. Plant Pathol.* **16**: 224-37.

Bal, S. and Pal, S. (2021) Seed Production Technology of Onion. *In: Text Book on Vegetable Seed Production*, Mishra, N., Panigrahi, K.K. and Mani, A. (Eds.). International Books & Periodical Supply Service, New Delhi, India, pp. 303-25.

Baldwin, S., Joyce, M.P., Wright, K., Chen, L. and McCallum, J. (2012a). Development of robust genomic simple sequence repeat markers for estimation of genetic diversity within and among bulb onion (*Allium cepa* L.) populations. *Mol. Breed.* **30**: 1401-11.

Baldwin, S., Revanna, R., Thomson, S., Pither-Joyc, Wright, K., Crowhurst, R., Fiers, M., Chen, L., Macknight, R. and McCallum, J.A. (2012b). A Toolkit for bulk PCR-based marker design from next-generation sequence data: application for development of a framework linkage map in bulb onion (*Allium cepa* L.). *BMC Genomics* **13**: 637

www.indianjournals.com
Not for Commercial Use
Downloaded From IP - 216.73.216.217 on dated 02-Nov-2026

Bark, O.H. and Havey, M.J. (1995). Similarities and relationships among populations of the bulb onion as estimated by nuclear RFLPs. *Theoret. Appl. Genetics* 90: 407-14. https://doi.org/10.1007/BF00221983

Berninger, E. (1965) Contribution a`l'e´tude de la sterilite´ ma^le de l'oignon (*Allium cepa* L.). *Ann. Ame´lior. Plant* **15**: 183-99.

Bohanec, B. (2002). Doubled-haploid Onions. *In: Allium crop science: recent advances,* Rabinowitch, H.D and Currah, L. (Eds.). Wallingford, UK: CABI Publishing, pp.145-57.

Bohanec, B. (2009). Double haploids. In: *Advances in Haploid Production in Higher Plants*, Touraev, A., Forster, B.P and Jain S.M. (Eds.). Springer, Heidelberg, pp. 35-46.

Bradeen, J.M. and Havey, M.J. (1995). Randomly amplified polymorphic DNA in bulb onion and its use to assess inbred integrity. *J. Am. Soc. Hort. Sci.* **120**: 752-58.

Brewster, J.L. (2008). *Onions and Other Vegetable Alliums*. CABI, Oxfordshire, UK.

Caldwell, T.J., Lada, R. and Hooper, D. (2003). Physiological responses of onion (*Allium cepa* L.) seedlings exposed to drought. *Acta Hortic*. **618**: 321-27.

Chand, S.K., Nanda, S. and Joshi, R.K. (2018). Genetics and molecular mapping of a novel purple blotch-resistant gene ApR1 in onion (*Allium cepa* L.) using STS and SSR markers. *Mol. Breed.* **38**:109.

Clarke, A.E., Jones, H.A. and Little, T.M. (1944). Inheritance of bulb colour in the onion. *Genetics* **29**: 569-75.

Cramer, C.S. (2000). Breeding and genetics of *Fusarium* basal rot resistance in onion. *Euphytica.* **115**: 159-66.

Damon, S.J. and Havey, M.J. (2014). Quantitative trait loci controlling amounts and types of epicuticular waxes in onion. *J. Amer. Soc. Hort. Sci.* **139**(5): 597-602.

Diaz-Montano, J., Onatno, J., Fuchs, M., Nault, B.A., Fail, J. and Shelton, A.M. (2011). Onion thrips (Thysanoptera: Thripidae): a global pest of increasing concern in onion. *J. Econ. Entomol.* **104**: 1-13.

Duangjit, J., Bohanec, B., Chan, A.P., Town, C.D. and Havey, M.J. (2013). Transcriptome sequencing to produce SNP-based genetic maps of onion. *Theor. Appl. Genet.* **126**: 2093-2101.

Eady, C.C., Kamoi, T., Kato, M., Porter, N.G., Davis, S., Shaw, M., Kamoi, A. and Imai, S. (2008). Silencing onion lachrymatory factor synthase causes a significant change in the sulfur secondary metabolite profile. *Plant Physiology* **147**(4): 2096-106.

Eidlin, Y.T., Monakhos, G.F. and Monakhos, S.G. (2021). Marker-assisted breeding of onion (*A. cepa* L.) maintainer line resistant to Downy mildew. *Vegetable Crops of Russia* **3**: 34-39.

El-Shafie, M.W. and Davis, G.N. (1967). Inheritance of bulb colour in onion (*Allium cepa* L.). *Hilgardia.* **38**(17): 607-22.

Esfahani, M.N. and Pour, B.A. (2008). Differences in resistance in onion cultivars to pink root rot disease in Iran. J. Gen. *Plant Pathol.* **74**: 46-52.

FAO STAT (2024). Area production and productivity of dry onion in the world. Food and Agricultural Organization of the United Nations, Rome, Italy. Available at: http://www.fao.org/faostat/en/#data/QC. Date assessed: 03/03/2026.

Galmarini, C.R., Goldman, I.L. and Havey, M.J. (2001). Genetic analyses of correlated solids, flavor, and health-enhancing traits in onion (*Allium cepa* L.). *Mol. Genet. Genomics*. **265**: 543-51.

Galvan, G.A., Carole, F.S., Boucoiran, K., Koopman, W.J.M., Meijer, K.B., Gonzalez, P.H., Waalwijk, C., Kik, C. and Scholten, O.E. (2008). Genetic variation among Fusarium isolates from onion, and resistance to *Fusarium* basal rot in related *Allium* species. *Eur. J. Plant Pathol.* **121**: 499-512.

Glaszmann, J.C., Kilian, B., Upadhyaya, H.D. and Varshney, R.K. (2010). Accessing genetic diversity for crop improvement. *Curr. Opin. Plant Biol*. **13**: 1-7.

Go¨kce, A.F., McCallum, J., Sato, Y. and Havey, M.J. (2002). Molecular tagging of the *Ms* locus in onion. *J. Am. Soc. Hort. Sci.* **127**: 576-82.

Goldman, I.L. (2011). Molecular breeding of healthy vegetables. *EMBO Rep*. **12**: 96-102.

Gupta, A.J., Anandhan, S., Manjunathagowda, D.C., Benke, A.P., Mahajan, V., Kad, S.K. and Singh, M. (2022). Complement test for distinctiveness, uniformity and stability testing of *kharif* onion (*Allium cepa* L.) varieties. *Genetic Resources and Crop Evolution* **69**(6): 2217-29.

Havey, M.J., Galmarini, C.R., Gokce, A.F. and Henson, C. (2004). QTL affecting soluble carbohydrate concentrations in stored onion bulbs and their association with flavor and health-enhancing attributes. *Genome* **47**: 463-68.

Hovius, M.H.Y., Yandell, B. and Goldman, I.L. (2004). Evaluation of long-day onions for resistance to white rot infection using greenhouse and laboratory techniques. *J. Am. Soc. Hortic. Sci.* **129**: 258-65.

Huo, Y. M., Liu, B.J., Yang, Y.Y. and Wu, X. (2015). AcSKP1, a multiplex PCR-based co-dominant marker in complete linkage disequilibrium with the male-fertility restoration (*Ms*) locus, and its application in open pollinated populations of onion. *Euphytica* **204**(3): 711-22.

Insani, E.M., Cavagnaro, P.F., Salomón, V.M., Langman, L., Sance, M., Pazos, A.A., Carrari, F.O., Olga, Filippini, O., Vignera, L. and Galmarini, C.R. (2016). Variation for Health-Enhancing Compounds and Traits in Onion (*Allium cepa* L.) Germplasm. *Food and Nutrition Sciences* **7**: 577-91.

Ivchenko, T., Volkova, N., Bashtan, N., Korniyenko, S., Kurutko, R. and Ka, O.B. (2017). Identification of Onion

Downloaded From IP - 216.73.216.217 on dated 02-Nov-2026
Not for Commercial Use
www.indianjournals.com

Varieties (*Allium cepa* L.) of Ukrainian Breeding Using Microsatellite Markers. *Journal of Pharmaceutical, Chemical and Biological Sciences* **5**(1): 40-44.

Jakše, J., Meyer, J.D.F., Suzuki, G., McCallum, J., Cheung, F., Town, C.D. and Havey, M.J. (2008). Pilot sequencing of onion genomic DNA reveals fragments of transposable elements, low gene densities, and significant gene enrichment after methyl filtration. *Mol. Genet. Genomics* **280**: 287-92.

Jayaswall, K., Kumar, D., Jayaswall, D., Sagar, R., Kumar, S., Sharma, R.K., Bhagat, K., Mahajan, V., Sharma, H., Singh, S.P. and Sharma, I. (2025). Immunomodulator and marker-assisted identification of *Allium* genotypes containing immunological memory against anthracnose. *Physiological and Molecular Plant Pathology* **139**: 102815.

Jeon, S., Han, J., Kim, C.W., Kim, J.G., Moon, J.H. and Kim, S. (2022). Identification of a candidate gene responsible for the G locus determining chartreuse bulb colour in onion (*Allium cepa* L.) using bulked segregant RNA-Seq. *Theoretical and Applied Genetics* **135**(3): 1025-36.

Jo, J., Purushotham, P.M., Han, K., Lee, H.R., Nah, G. and Kang, B.C. (2017). Development of a Genetic Map for Onion (*Allium cepa* L.) Using Reference-Free Genotyping-by-Sequencing and SNP Assays. *Frontiers in Plant Science* 8(1606): 1-8.

Jones, H.A. and Clarke, A. (1943). Inheritance of male sterility in the onion and the production of hybrid seed. *Proc. Am. Soc. Hort. Sci.* **43**:189-94.

Jones, H.A. and Emsweller, S.L. (1936). A male-sterile onion. *Proc. Am. Soc. Hort. Sci.* **34**: 582-85.

Kamal, N., Saxena, A., Steiner, S.L. and Cramer, C.S. (2012). Screening of new Mexico autumn-sown onions for black mold disease. *Hort. Technology* **22**: 719-23.

Kareem, M.A., Murthy, K.V.M.K., Nadaf, H.A. and Waseem, M.A. (2012). Effect of temperature, relative humidity and light on lesion length due to *Alternaria porri* in onion. *Asian J. Environ. Sci.* **7:** 47-49.

Kato, M., Masamura, N., Shono, J., Okamato, D., Abe, T. and Imai, S. (2016). Production and characterization of tearless and non-pungent onion. *Scientific Reports* **6**: 23779.

Kesraliker, M., Mokate, S., Golakiya, B.A. and Vakharia, D.N. (2017). Genetic Diversity Analysis among Onion Genotypes through RAPD, SSR and ISSR markers. *Int. J. Curr. Microbiol. App. Sci.* **6**(6): 1467-74.

Khar, A., Jaksey, J. and Havey, M.J. (2008). Segregations for onion bulb colors reveal that red is controlled by at least three loci. *J. Amer. Soc. Hort. Sci.* **133**(1): 42-47.

Khar, A., Zimik, M., Verma, P., Singh, H., Mangal, M., Singh, M.C. and Gupta, A.J. (2022). Molecular marker-based characterization of cytoplasm and restorer of male sterility (*Ms*) locus in commercially grown onions in India. *Mol. Biol. Rep*. **49**(6): 5535-45.

Khosa, J.S., McCallum, J., Dhatt, A.S. and Macknight, R.C. (2016). Enhancing onion breeding using molecular tools. *Plant Breeding* **135**: 9-20.

Khosa, J.S., Negi, K.S., Dhatt, A.S. and Khar, A. (2014). Utility of Simple Sequence Repeat (SSR) markers to realize worth of germplasm in genus *Allium*. *Indian J. Plant Genet. Resour.* **27**: 238-45.

Kim, B. and Kim, S. (2019). Identification of a variant of CMS-T cytoplasm and development of high resolution melting markers for distinguishing cytoplasm types and genotyping a restorer-of-fertility locus in onion (*Allium cepa* L.). *Euphytica* **215**: 164.

Kim, S. (2014). A codominant molecular marker in linkage disequilibrium with a restorer-of-fertility gene (*Ms*) and its application in re-evaluation of inheritance of fertility restoration in onions. *Mol. Breed.* **34**: 769-78. https://doi.org/10.1007/s11032-014-0073-8

Kim, S., Jones, R., Yoo, K. and Pike, L.M. (2004). Gold colour in onions (*A. cepa*): a natural mutation of the *chalcone isomerase* gene resulting in a premature termination codon. *Molecular Genetics and Genomics* **272**: 411-19.

Kim, S., Kim, C.W., Park, M. and Choi, D. (2015). Identification of candidate genes associated with fertility restoration of cytoplasmic male-sterility in onion (*Allium cepa* L.) using a combination of bulked segregant analysis and RNA-seq *Theor. Appl. Genet.* **128**: 2289- 99.

Kim, S.J., Park, J.S., Park, T., Lee, H.M., Choi, J.R. and Park, Y.D. (2021). Development of molecular markers associated with resistance to gray mold disease in onion (*Allium cepa* L.) through RAPD-PCR and transcriptome analysis. *Horticulturae* **7**(11): 436.

King, J.J., Bradeen, J.M., Bark, O., McCallum, J.A. and Havey, M.J. (1998). A low-density genetic map of onion reveals a role for tandem duplication in the evolution of an extremely large diploid genome. *Theor. Appl. Genet.* **96**: 52-62.

Kuhl, J.C., Cheung, F., Yuan, Q.P., Martin, W., Zewdie, Y., McCallum, J., Catanach, A., Rutherford, P., Sink, K.C., Jenderek, M., et al. (2004). A unique set of 11,008 onion expressed sequence tags reveals expressed sequence and genomic differences between the monocot orders Asparagales and Poales. *Plant Cell* **16**: 114-25.

Labate, J.A., Glaubitz, J.C. and Havey, M.J. (2020). Genotyping by sequencing for SNP marker development in onion. *Genome* **63**(12): 607-13.

Lee, Y.R., Kim, C.W., Han, J., Choi, H.J., Han, K., Lee, E.S., Kim, D.S., Lee, J., Siddique, M.I. and Lee, H.E. (2021). Genotyping by sequencing derived genetic linkage map and quantitative trait loci for sugar content in onion (*Allium cepa* L.). *Plants* 10(11): 2267.

Litt, M. and Luty, J.A. (1989). A hypervariable microsatellite revealed by in vitro amplification of a dinucleotide

www.indianjournals.com
Not for Commercial Use
Downloaded From IP - 216.73.216.217 on dated 02-Nov-2026

repeat within the cardiac muscle actin gene. *American Journal of Human Genetics* **44**: 397-401.

Mahala, K., Yadav, D.K., Gupta, D., Choudhary, M.R., Ujjainiya, P. and Bhateshwar, M.C. (2024). Estimation of magnitude of heterosis for growth traits in onion (*Allium cepa*). *Current Horticulture* **12**(3): 68-70.

Mainkar, P., Manape, T.K., Satheesh, V. and Anandhan, S. (2023). CRISPR/Cas9-mediated editing of PHYTOENE DESATURASE gene in onion (*Allium cepa* L.). *Frontiers in Plant Science* **14**: 1226911.

Mallor, C., Balcells, M., Mallor and Sales, E. (2011). Genetic variation for bulb size, soluble solids content and pungency in the Spanish sweet onion variety Fuentes de Ebro Response to selection for low pungency. *Plant Breeding* **130**: 55-59.

Maniruzzaman, M., Haque, M.E., Haque, M.M., Sayem, M.A. and Al-Amin, M. (2010). Molecular characterization of onion (*Allium cepa*) using RAPD markers. *Bangladesh Journal of Agricultural Research* **35**(2): 313-22.

Manjunathagowda, D.C. and Selvakumar, R. (2021). Marker-assisted selection of *Ms* locus responsible for male fertility restoration in onion (*Allium cepa* L.). *Genetic Resources and Crop Evolution* **68**: 2793-97.

Manjunathagowda, D.C. (2025). Marker assisted selection of male sterile (CMS-S) lines among short day Indian onion genotypes. *Vegetos* **38**(5): 1906-10.

Martin, W., McCallum, J., Shigyo, M., Jakse, J., Kuhl, J.C., Yamane, N., Sink, K.C., Town, C.D and Havey, M.J. (2005). Genetic mapping of expressed sequences in onion and in silico comparisons show scant colinearity with rice. *Mol. Genet. Genom.* **274**: 197-204.

Masamura, N., McCallum, J., Khrustaleva, L., Kenel, F., Pither-Joyce, M., Shono, J., Suzuki, G., Yamauchi, N. and Shigyo, M. (2012). Chromosomal organization and sequence diversity of genes encoding lachrymatory factor synthase in *Allium cepa* L. *Genes Genomes Genomics* **2**: 643-51.

McCallum, J. (2007). Onion. In: *Genome Mapping and Molecular Breeding in Plants*. Volume 5 Vegetables, Kole, C. (Ed.). Springer, Berlin Heidelberg, pp. 331-347.

McCallum, J., Andrew, C., Meeghan, P.J., Martin, S., Ruth, B., John, S., Ian, S., Masayoshi, S. and Havey, M.J. (2006). Genetic mapping of a major gene affecting onion bulb fructan content. *Theor. Appl. Genet.* **112**: 958-67.

McCallum, J., Baldwin, S., Shigyo, M., Deng, Y., Heusden, S.V., Pither-Joyce, M. and Kenel, F. (2012). *Allium*Map-A comparative genomics resource for cultivated *Allium* vegetables. *BMC Genomics* **13**: 168.

McCallum, J., Leite, D., Pither-Joyce, M. and Havey, M.J. (2001). Expressed sequence markers for genetic analysis of bulb onion (*Allium cepa* L.). *Theor. Appl. Genet.* **103**: 979-91.

McCallum, J., Pither-Joyce, M., Shaw, M., Kenel, F., Davis, S., Butler, R., Scheffer, J., Jakse, J. and Havey, M.J. (2007). Genetic mapping of sulphur assimilation genes reveals a QTL for onion bulb pungency. *Theor. Appl. Genet.* **114**: 815-22.

McCallum, J., Thomson, S., Pitcher-Joyce, M., Kenel, F., Clarke, A. and Havey, M.J. (2008). Genetic diversity analysis and single nucleotide polymorphism marker development in cultivated bulb onion based on expressed sequence Tag-simple sequence repeats markers. *J. Am. Soc. Hortic. Sci.* **113**: 810-18.

Mitrová, K., Svoboda, P. and Ovesná, J. (2015). The selection and validation of a marker set for the differentiation of onion cultivars from the Czech Republic. *Czech J. Genet. Plant Breed.* 51: 62-67.

Mukherjee, A., Sikdar, B., Ghosh, B., Banerjee, A., Ghosh, E., Bhattacharya, M. and Roy, S.C. (2013). RAPD and ISSR analysis of some economically important species, varieties, and cultivars of the genus *Allium* (Alliaceae). *Turkish Journal of Botany* **37**: 605-18.

Nanda, S., Chand, S.K., Mandal, P., Tripathi, P. and Joshi, R.K. (2016). Identification of novel source of resistance and differential response of Allium genotypes to purple blotch pathogen, *Alternaria porri* (Ellis). *Ciferri. Plant Pathol. J.* **32**(6): 519-27.

Ohara, T., Song, Y.S., Tsukazaki, W., Nunome, T. and Kojima, A. (2005). Genetic mapping of AFLP markers in Japanese bunching onion (*Alliums fistulosum*). *Euphytica* **144**: 255-63.

Pal, S., Krishna, R., Dedhia, L., Panwar, H.S., Karkute, S.G., Rai, N., Kumar, R., Pandey, S. and Singh, A.K. (2025). CRISPR mediated gene editing for economically important traits in horticultural crops: progress and prospects. *Transgenic Research* **34**: 26.

Pal, S., Revadi, M., Thontadarya, R.N., Reddy, D.C.L., Varalakshmi, B., Pandey, C.D. and Rao, E.S. (2020). Understanding genetic diversity, population structure and development of a core collection of Indian accessions of watermelon (*Citrullus lanatus* (Thunb.) Matsum. and Nakai). *Plant Genetic Resources: Characterization and Utilization* **18**(5): 359-68.

Park, J., Cho, D.Y., Moon, J.S., Yoon, M.K. and Kim, S. (2013). Development of functional markers for detection of inactive *DFR-A* alleles responsible for failure of anthocyanin production in onions (*Allium cepa* L.). *Korean Journal of Horticultural Science and Technology* **31**: 72-79.

Priya, R.U., Sataraddi, A. and Darshan, S. (2016). Survey for purple blotch [*Alternari porri* (Ellis) Cif.] on onion (*Allium cepa*) in northern parts of Karnataka. *Current Horticulture* **4**(1): 55-60.

Raines, S., Henson, C. and Havey, M.J. (2009). Genetic analyses of soluble carbohydrate concentration in onion bulbs. *Journal of the American Society for Horticultural Science* **134**: 618-23.

Downloaded From IP - 216.73.216.217 on dated 02-Nov-2026
Not for Commercial Use
www.indianjournals.com

Sahoo, J., Mahanty, B., Mishra, R. and Joshi, R.K. (2023). Development of SNP markers linked to purple blotch resistance for marker-assisted selection in onion (*Allium cepa* L.) breeding. *3 Biotech* **13**(5): 137.

Saini, N., Hedau, N.K., Khar, A., Yadav, S., Bhatt, J.C. and Agarwal, P.K. (2015). Successful deployment of marker assisted selection (MAS) for inbred and hybrid development in long-day onion (*Allium cepa* L.). *Indian Journal of Genetics* **75**(1): 93-98.

Sato, Y. (1998). PCR amplification of 671 CMS-specific mitochondrial nucleotide sequences to identify cytoplasmic genotypes of onion (*Allium cepa* L.). *Theoretical and Applied Genetics* **96**: 367-70.

Scholten, O.E., Heusden, A.W.V., Khrustaleva, L.I., Burger-Meijer, K., Mank, R.A., Antonise, R.G.C., Harrewijn, J.L., Van-Haecke, W., Oost, E.H., Peters, R.J. and Kik, C. (2007). The long and winding road leading to the successful introgression of downy mildew resistance into onion. *Euphytica* **156**: 345-53.

Scholten, O.E., van Kaauwen, M.P. and Shahin, A. (2016). SNP-markers in *Allium* species to facilitate introgression breeding in onion. *BMC Plant Biology* **16**: 187.

Schwartz, H.F., Gent, D.H. and Bartolo, M.E. (2005). Purple blotch. http://wwwhighplainsipm.org/HpIPMSearch/docs/purple blotch-onion.htm

Schweisguth, B. (1973). E´tude d'un nouveau type de ste´rilite´ male chez l'oignon, *Allium cepa* L. *Ann. Ame´lior. Plant* **23**: 221-33.

Schwinn, K.E., Ngo, H., Kenel, F., Brummell, D.A., Albert, N.W., McCallum, J.A., Pither-Joyce, Crowhurst, R.N., Eady. C. and Davies, K.M. (2016). The onion (*Allium cepa* L.) R2R3-MYB gene regulates anthocyanin biosynthesis. *Frontiers in Plant Science* **7**: 1865.

Singh, H., Khar, A. and Verma, P. (2021), Induced mutagenesis for genetic improvement of *Allium* genetic resources: a comprehensive review. *Genetic Resources and Crop Evolution* **1**: 461-68.

Singh, H.P. (2010). Ongoing research in abiotic stress due to climate change in horticulture, Curtain Raiser Meet on Research Needs Arising due to Abiotic Stresses in Agriculture Management in India under Global Climate Change Scenario, Baramati, Maharashtra, October 29-30. pp. 1-23. (http://www.niam.res.in/pdfs/DDG_Hort_lecture.pdf).

Slimestad, R., Fossen, T. and Vågen, I.M. (2007). Onions: a source of unique dietary flavonoids. *Journal of Agricultural Food Chemistry* **55**: 10067-80.

Smolik, M., Rzepka-Plevneš, R., Kowalczys, K. and Grabiec, M. (2007). Study of genetic diversity of the onion species by ISSR-PCR analysis. *Biotechnologia* **6**(1): 13-21.

Sudha, G.S., Ramesh, P., Sekhar, A.C., Krishna, T.S., Bramhachari, P.V. and Riazunnisa, K. (2019). Genetic diversity analysis of selected onion (*Allium cepa* L.) germplasm using specific RAPD and ISSR polymorphism markers. *Biocatalysis and Agricultural Biotechnology* **17**: 110-18.

Taylor, A.V., Vagany, D.J., Barbara, B., Thomas, D.A.C., Pink, J.E., Jones and Clarkson, J.P. (2013). Identification of differential resistance to six *Fusarium oxysporum* f. sp. *cepae* isolates in commercial onion cultivars through the development of a rapid seedling assay. *Plant. Pathol.* **62**: 103-11.

Tripathi, P., Priyadarshini, A., Das, S.K., Sahoo, B.B. and Dash, D.K. (2013). Evaluation of onion (*Allium cepa* L.) genotypes for tolerance to thrips (*Thrips tabaci* L.) and purple blotch (*Alternaria porri* (E) Ciferri). *Int. J. Biores. Stress Manage.* **4**: 561-64.

Tripathi, P.C. and Lawande, K.E. (2019) Onion storage in tropical region- a review. *Current Horticulture* **7**(2): 15-27.

Tripathi, P.C., Shankar, V. and Lawande, K.E. (2017) Microirrigation in onion (*Allium cepa*) and garlic (A. sativum)-a review. Current Horticulture **5**(1): 3-14.

Tsukazaki, H., Nunome, T., Fukuoka, H., Kanamori, H., Kono, I., Ohara, T., et al. (2008). Applications of DNA marker technology in Japanese bunching onion breeding. *Acta Horticulturae* **770**: 153-58.

Tsukazaki, H., Yaguchi, S., Yamashita, K., Kojima, A. and Wako, T. (2012). QTL mapping for bulb pigmentation in F_2 population between bulb onion and shallot. *Acta Horticulturae* **969**: 85-89.

Van Heusden, A.W., van-Ooijen, J.W., Vrielink-van Ginkel, R., Verbeek, W.H.J., Wietsma, W.A. and Kik, C. (2000). A genetic map of an interspecific cross in *Allium* based on amplified fragment length polymorphism (AFLPTM) markers. *Theor. Appl. Genet.* **100**: 118-26.

Varshney, R.K., Ribaut, J.M., Buckler, E.S., Tuberosa, R., Rafalski, J.A. and Langridge, P. (2012). Can genomics boost productivity of orphan crops? *Nat. Biotechnol.* **30**: 1172-76.

Yaguchi, S., McCallum, J., Shaw, M., Pither-Joyce, M., Onodera, S., Shiomi, N., Yamauchi, N. and Shigyo, M. (2008). Biochemical and Genetic Analysis of Carbohydrate Accumulation in *Allium cepa* L. *Plant Cell Physiol.* **49**(5): 730-39.

Zabeau, M. and Vos, P. (1993). Selective restriction fragment amplification: A general method for DNA fingerprinting. European patent application no. 92402629.7, publication no. 0534858 A1

Ziętkiewicz, E., Rafalski, A. and Labuda, D. (1994). Genome fingerprinting by simple sequence repeat (SSR)-anchored polymerase chain reaction amplification. *Genomics* **20**: 176-83.