Soil Functional Diversity in Selected Rainfed Groundnut Production Systems of India

M. Manjunath*[1], G. Ravindra Chary[1], K.C. Nataraja[2], B. Sahadeva Reddy[2], H.S. Latha[3], Mudalagiriappa[3], N.I. Patel[4], D.S. Hirpara[5], N. Jyothi lakshmi[1], A.K. Indoria[1], G. Pratibha[1], K.A. Gopinath[1], S. Savitha[1], Arun Shanker[1], M. Srinivasa Rao[1] and V.K. Singh[1]

[1]ICAR-Central Research Institute for Dryland Agriculture, Hyderabad–500 059, Telangana

[2]AICRP for Dryland Agriculture, ANGRAU, Ananthapuramu-515 001, Andhra Pradesh

[3]AICRP for Dryland Agriculture, UAS, Bengaluru-560 065, Karnataka

[3]AICRP for Dryland Agriculture, SDAU, Sardar Krishinagar-385 506, Gujarat

[5]AICRP for Dryland Agriculture, JAU, Rajkot-360 003, Gujarat

*E-mail: m.manjunath@icar.gov.in

ABSTRACT: Soil microorganisms are key drivers of various functions in agroecosystem. Soil dehydrogenase, alkaline and acid phosphatase enzymes are regarded as good indicators of microbiological activity of soils. The aim of the study was to assess the status of microbial activity in selected rainfed groundnut growing regions. Soil samples were collected from Bengaluru (Karnataka), Ananthapuramu (Andhra Pradesh), Sardar Krishinagar (Gujarat) and Rajkot (Gujarat) representing semi-arid (hot moist), arid (hot), semi-arid (hot dry), and semi-arid (hot dry) climatic conditions respectively. Microbial activities were estimated in terms of soil enzymes such as dehydrogenase, acid and alkaline phosphatase. The differences in microbial parameters across different regions were observed. Significant difference in dehydrogenase activity was observed between Sardar Krishinagar and Ananthapuramu soil samples. Higher soil functional diversity index was observed in Rajkot and Sardar Krishinagar. The results of the study suggest that, application of organic sources of nutrients like farm yard manure is essential to maintain soil microbial activity, functional diversity and soil health in rainfed groundnut production systems.

Keywords: Groundnut, microorganisms, rainfed, soil enzymes, soil health

www.IndianJournals.com
Not for Commercial Use
Downloaded From IP - 216.73.216.146 on dated 15-Nov-2026

Introduction

Soil microorganisms are indispensable part of agroecosystem as they perform multiple functions viz., organic matter decomposition, biogeochemical cycling of plant nutrients, mobilization, solubilization and transformation of nutrients, pest and disease management (Suman *et al.,*2022; Manjunath *et al.,* 2016 and 2017; Pratibha *et al.,* 2023). Soil enzymes are largely produced by microorganisms. They are vital for mineralization of soil organic matter, nutrient recycling and removal of toxic pollutants (Sainju *et al.,* 2022). They are largely used as indicators of microbial activity and soil health. Soil dehydrogenase, alkaline and acid phosphatase enzymes are regarded as good indicators of soil microbiological activity (Garcia *et al.,* 1997; Symanowicz *et al.,* 2022). Activities of soil micro-organisms are influenced by regional differences in climate, soil type and hydrothermal situations Srinivasarao *et al.,* 2018; Rampant use of inorganic fertilizers affects the diversity of soil microorganisms (Bharathi *et al.,* 2024). Decrease in soil microbial diversity would negatively impact the fertility and productivity of soil (Onet *et al.,* 2016). Groundnut is a major oilseed crop largely grown under rainfed conditions in India. Limited information is available on the status of soil microbial activity in different rainfed groundnut growing regions of India. Hence, the aim of the present study was to assess the status of microbial activity in selected rainfed groundnut growing regions viz., Bengaluru (Karnataka), Ananthapuramu (Andhra Pradesh), Rajkot (Gujarat) and Sardar Krishinagar (SK Nagar, Gujarat) representing semi-arid (hot moist), arid (hot), semi-arid (hot dry) and semi-arid (hot dry) climatic conditions respectively.

Materials and methods

Study area and collection of soil samples

The study area consisted of major rainfed groundnut growing regions viz., Ananthapuramu, Bengaluru, Sardar Krishinagar and Rajkot. The climatic and edaphic features of the studied rainfed groundnut growing locations have been presented in Table 1. Soil samples were collected from five farmer's field each from nearby areas of Ananthapuramu, Bengaluru, Sardar Krishinagar and Rajkot AICRPDA centers. At each farmer's field, after harvest of crop, four random samples (0-15 cm depth) were collected and pooled. The soil samples were immediately kept in cool boxes after removal of root pieces and gravel. The soil samples were brought to laboratory and stored at 4°C for further analysis.

Table 1: Climatic and edaphic characteristics of the studied locations

AICRPDA Centre	Agroclimatic Zone	Climate	Soil type	Rainfall (mm)
Ananthapu-ramu	Scarce rainfall zone (Rayalaseema) in Andhra Pradesh (3.0)	Arid (Hot)	Alfisols	544
Rajkot	North Saurashtra zones in Gujarat (5.1)	Semiarid (Hot dry)	Vertisols	590
SK Nagar	Northern Gujarat in Gujarat (2.3)	Semiarid/ Arid (Hot dry)	Entisols	670
Bengaluru	Central, eastern and southern dry zone in Karnataka (8.2)	Semiarid (Hot moist)	Alfisols	926

www.IndianJournals.com
Not for Commercial Use
Downloaded From IP - 216.73.216.146 on dated 15-Nov-2026

Analysis of microbial parameters

Soil dehydrogenase activity was measured spectrophotometrically using 2, 3, 5 triphenyl tetrazolium chloride (TTC) as substrate (Casida *et al.,* 1964). Alkaline and acid phosphatase enzyme activities were determined (Tabatabai and Bremner, 1969). Soil functional diversity was determined as per the procedure described by Rodriguez-Loinaz *et al.* (2008) and Kompała Bąba *et al.* (2021) with the following formula;

$$\text{Soil H}' = \sum_{i=1}^{s} pi \ln pi$$

where pi is the ratio of the activity of a specific enzyme to the sum of all the measured enzyme activities.


Statistical analysis

One way analysis of variance (ANOVA) of the data was done using SPSS 16.0 (SPSS Inc., Chicago, IL, USA) for windows. Comparison of means was made using the least significant difference at $p \leq 0.05$.

Results and Discussion

Soil dehydrogenase activity

Soil dehydrogenase activity ranged from 28.73 to 32.03 TPF µg/g/d in the studied rainfed groundnut growing regions viz., Ananthapuramu, Bengaluru, Sardar Krishinagar and Rajkot. Significantly ($p \leq 0.05$) higher dehydrogenase activity was recorded in SK Nagar (32.03 TPF µg/g/d), however it was at par with Bengaluru and Rajkot soil samples. Lowest dehydrogenase activity was observed in Ananthapuramu soil samples (Fig. 1).




(Note: Data presented is means ±SE. Different alphabets at each column indicates significant difference (p≤0.05))

Fig. 1: Soil dehydrogenase enzyme activity (TPF µg/g/d) at selected rainfed groundnut growing locations

Alkaline phosphatase activity

Relatively higher and statistically ($p \leq 0.05$) at par values of alkaline phosphatase activity was observed in SK Nagar and Rajkot soil samples. Lowest alkaline phosphatase activity (7.25 PNP ug/g/h) was observed in Bengaluru soil samples (Fig.2).




(Note: Data presented is means±SE*. Different alphabets at each column indicates significant difference (p≤0.05))*

Fig. 2: Alkaline phosphatase enzyme activity (PNP ug/g/h) at selected rainfed groundnut growing locations

Acid phosphatase activity

Maximum acid phosphatase activity (10.76 PNP ug/g/h) was recorded in Bengaluru soil samples followed by Rajkot. Lowest acid phosphatase activity (8.09 PNP ug/g/h) was noticed in Ananthapuramu soil samples, however it was statistically at par with SK Nagar and Rajkot soil samples (Fig. 3).




(Note: Data presented is means ±SE. Different alphabets at each column indicates significant difference (p≤0.05))

Fig. 3: Acid phosphatase enzyme activity at (PNP ug/g/h) selected rainfed groundnut growing locations

Soil functional diversity

Rajkot samples recorded significantly higher soil functional diversity index (H'=1.099) as compared to Ananthapuramu and Bengaluru soil samples. No significant difference in soil functional diversity index was observed between Bengaluru and Ananthapuramu soil samples (Fig. 4).



(Note: Data presented is means ±SE. Different alphabets at each column indicates significant difference (p≤0.05))

Fig. 4: Soil functional diversity index of selected rainfed groundnut growing locations

Soil microorganisms are the major source of soil enzymes (Bandick and Dick, 1999). Soil enzymes are very important for organic matter decomposition and cycling of various nutrients **(**Uwituze *et al.,* 2022). They act as early indicators of soil health (Sainju *et al.,* 2022) and have been widely used to monitor the soil health (Lee *et al.,* 2020). Dehydrogenase is an intracellular enzyme that functions only in living microorganisms hence, it is estimated to know the activity of living or viable microorganisms in the soil. In the present study, statistically at par values for dehydrogenase activity was observed in SK Nagar, Bengaluru and Rajkot soil samples. It could be due to regular use of FYM in the fields. Addition of FYM stimulates soil microorganisms due to the provision of organic carbon and other nutrients in easily available form (Kumar *et al.,* 2021). Soil treated with FYM and organic manure showed higher levels of dehydrogenase activity as compared to soil treated with mineral fertilizers (Lee *et al.,* 2004), reported that, incorporation of FYM enhanced the soil dehydrogenase activity. Acid and alkaline phosphatase enzymes convert organic phosphorus into inorganic phosphorus and improve fertility of soil (Kalayu, 2019; Chaudhary *et al.,* 2022). In the present investigation, maximum alkaline phosphatase activity was noticed in SK Nagar and Rajkot soil samples. Whereas Bengaluru soil samples recorded higher acid phosphatase activity as compared to other centers. Soil microbial functional diversity is important for organic matter decomposition and cycling of nutrients (JezierskaTys *et al.,* 2020). Organic sources of nutrients improved the soil microbial diversity (Manjunath *et al.,* 2018). In the present study, relatively maximum soil functional diversity index was observed in Rajkot followed SK Nagar soil samples. Higher functional diversity implies greater capability of utilization of different substrates by the microorganisms and participation in various transformation processes in soil.

Conclusion

Overall, higher soil microbial activity and functional diversity was recorded in Rajkot and SK Nagar soil samples. Inclusion of organic sources of nutrients such as farmyard manure in rainfed groundnut production is important for improving soil microbial activity, functional diversity and soil health.

Acknowledgement

The authors thank the authorities of ICAR-CRIDA, AICRPDA, Hyderabad; AICRPDA center, SKNDAU, SK Nagar; AICRPDA center, UAS (B), Bengaluru; AICRPDA center, JAU, Rajkot and AICRPDA center, ANGRAU, Ananthapuramu for extending required facilities for taking up this study.

References

Bandick AK and Dick RP. 1999. Field management effects on soil enzyme activities. *Soil Biology and Biochemistry,* 31: 1471–1479

Bharathi JM, Anbarasu M, Ragu R and Subramanian E. 2024. Assessment of soil microbial diversity and soil enzyme activities under inorganic input sources on maize and rice ecosystems. *Saudi Journal of Biological Sciences,* 31(5): 103978

Casida LE, Jr Klein DA and Santoro T. 1964. Soil dehydrogenase activity. *Soil Science,* 98: 371–376

Chaudhary P, Chaudhary A, Bhatt P, Kumar G, Khatoon H, Rani A, Kumar S and Sharma A. 2022. Assessment of soil health indicators under the influence of nanocompounds and *Bacillus* spp. in field condition. *Frontiers in Environmental Science,* 9: 769871.

www.IndianJournals.com
Not for Commercial Use
Downloaded From IP - 216.73.216.146 on dated 15-Nov-2026

Garcia C, Hernández MT and Costa F. 1997. Potential use of dehydrogenase activity as an index of microbial activity in degraded soils. *Communications in Soil Science and Plant Analysis,* 28: 123–134

Jezierska-Tys S, Wesołowska S and Gałązka A. 2020. Biological activity and functional diversity in soil in different cultivation systems. *International Journal of Environmental Science and Technology,* 17: 4189–4204

Kalayu G. 2019. Phosphate solubilizing microorganisms: Promising approach as Biofertilizers. *International Journal of Agronomy,* 4917256

Kompała-Bąba A, Bierza W, Sierka E, Błońska A, Besenyei L and Woźniak G. 2021.The role of plants and soil properties in the enzyme activities of substrates on hard coal mine spoil heaps. *Scientific Reports,* 11(1): 5155.

Kumar S, Dhar S, Barthakur S, Rajawat MVS, Kochewad SA, Kumar S, Kumar D and Meena LR. 2021. Farmyard manure as k-fertilizer modulates soil biological activities and yield of wheat using the integrated fertilization approach. *Frontiers in Environmental Science,* 9: 764489.

Lee JJ, Park RD, Kim YW, Shim JH, Chae DH, Rim YS and Kyoon B. 2004. Effect of food waste compost on microbial population, soil enzyme activity and lettuce growth. *Journal of Bioresource Technology,* 93: 21-28

Lee SH, Kim MS, Kim JG and Kim SO. 2020. Use of soil enzymes as indicators for contaminated soil monitoring and sustainable management. *Sustainability*, 12(19): 8209.

Manjunath M, Kanchan A, Ranjan K, Venkatachalam S, Prasanna R, Ramakrishnan B, Hossain F, Nain L, Shivay YS, Rai AB and Singh B. 2016. Beneficial cyanobacteria and eubacteria synergistically enhance bioavailability of soil nutrients and yield of okra. *Heliyon,* 2(2): e00066

Manjunath M, Kumar U, Yadava R B, Rai AB and Singh B. 2018. Influence of organic and inorganic sources of nutrients on the functional diversity of microbial communities in the vegetable cropping system of the Indo-Gangetic plains. *Comptes Rendus Biologies,* 341(6): 349-357.

Manjunath M, Singh A, Tripathi AN, Prasanna R, Rai AB and Singh B. 2017. Bioprospecting the fungicides compatible *Trichoderma asperellum* isolate effective against multiple plant pathogens in vitro. *Journal of Environmental Biology,* 38(4): 553.

Manjunath M, Yadav SK, Khokhar A, Patil SB, Asewar BV, Babu VS, Gopinath KA, Chary GR, Jyothilakshmi N, Srinivas K, Prabhakar M and Singh VK. 2022. Diversity of culturable soil microorganisms in different rainfed regions of India. *Environment and Ecology,* 40 (2A): 528-535

Onet A, Cimpeanu C, Teusdea A, Pantea S and Modog T. 2016. Evaluation of the soil properties variability in relation to different crop types. *Journal of Environmental Protection and Ecology,* 17:1305–1314.

Pratibha G, Manjunath M, Raju BMK, Srinivas I, Rao KV, Shanker AK, Prasad JVNS, Rao MS, Kundu S, Indoria AK, Kumar U, Rao KS, Anna S, Rao CS, Singh VK, Biswas AK and Chaudhari SK. 2023. Soil bacterial community structure and functioning in a long-term conservation agriculture experiment under semi-arid rainfed production system. *Frontiers in Microbiology,* 14: 1102682.

Rodriguez-Loinaz G, Onaindia M, Amezaga I, Mijangos I and Garbisu C. 2008. Relationship between vegetation diversity and soil functional diversity in native mixed-oak forests. *Soil Biology and Biochemistry,* 40: 49–60.

Sainju UM, Liptzin D and Dangi SM. 2022. Enzyme activities as soil health indicators in relation to soil characteristics and crop production. *Agrosystems, Geosciences and Environment,* 5: e20297

Srinivasarao Ch, Kundu S, Grover M, Manjunath M, Sudhanshu KS, Patel JJ, Singh SR, Singh RP, Patel MM, Arunachalam A and Soam SK. 2018. Soil enzymes in semi-arid and sub-humid rainfed agricultural systems as indicators of soil sustainability. Tropical Ecology. 59(1): 99-108.

Suman J, Rakshit A, Ogireddy SD, Singh S, Gupta C and Chandrakala J. 2022. Microbiome as a key player in sustainable agriculture and human health. *Frontiers in Soil Science,* 2: 821589.

Symanowicz B, Toczko R and Toczko, M. 2022. Enzymatic activity of soil after applying mineral fertilizers and waste lignite to maize grown for silage. *Agriculture,* 12(12): 2146.

Tabatabai MA and Bremner JM. 1969. Use of p-nitrophenyl phosphate for assay of soil phosphatase activity. *Soil Biology and Biochemistry,* 1: 301–307.

Uwituze Y, Nyiraneza J, Fraser TD, Dessureaut-Rompré J, Ziadi N and Lafond J. 2022. Carbon, nitrogen, phosphorus and extracellular soil enzyme responses to different land use. *Frontiers in Soil Science,* 2: 814554.

Srinivasarao Ch, Kundu S, Grover M, Manjunath M, Sudhanshu KS, Patel JJ, Singh SR, Singh RP, Patel MM, Arunachalam A and Soam SK. 2018. Soil Enzymes in semi-arid and sub-humid rainfed agricultural systems as indicators of soil sustainability. Tropical Ecology, 59(1): 99-108.

www.IndianJournals.com
Not for Commercial Use
Downloaded From IP - 216.73.216.146 on dated 15-Nov-2026

Received: March 2024 Accepted: May 2024