

Review

Lumpy skin disease (LSD): a risk to farmers and livestock worldwide

ANJU BALA¹, UMESH KUMAR JAISWAL¹, ANJU POONIA^{2*}, ANAMIKA³, ANIKA MALIK¹
and RACHNA⁴

¹Department of Veterinary and Animal Husbandry Extension Education

²Department of Veterinary Surgery and Radiology, ³Department of Animal Genetics and Breeding

⁴Department of Dairy Business Management

College of Veterinary Sciences, Lala Lajpat Rai University of Veterinary and Animal Sciences
Hisar 125004 Haryana, India

*Email for correspondence: anjupooniavs@gmail.com

© Society for Advancement of Human and Nature (SADHNA)

Received: 17.04.2023/Accepted: 24.05.2023

ABSTRACT

Lumpy skin disease is an emerging bovine viral disease which is endemic in most African countries and some Middle East ones and the elevated risk of the spread of disease into the rest of Asia and Europe should be considered. Understanding the restrictions and channels of distribution is crucial given the recent rapid spread of disease in nations that are currently disease-free. *Capripoxvirus*, the responsible agent, can also cause goatpox and sheepox. Given that they pose a threat to worldwide trade and could be exploited as tools of economic bio-terrorism, the economic impact of these diseases is a major worry. Due to poor conditions in farming communities and limited availability to efficient immunizations, the dissemination of *Capripoxvirus* appears to be spreading. This is mostly caused by the economic consequences of the Covid-19 pandemic, the enforcement of debilitating sanctions in endemic areas, a growth in the trading in live animals and animal products, both legally and illegally, as well as the repercussions of global climate change. The goal of the current review is to offer current knowledge on the different facets of the illness, including its clinicopathology, transmission, epidemiology, diagnostics, preventative and control strategies and the probable contribution of wildlife to the disease's spread.

Keywords: Capripox; disease; epidemiology; lumpy skin disease; livestock


INTRODUCTION

Lumpy skin disease (LSD) is an infectious viral disease of cattle and buffaloes caused by the *Capripoxvirus* of family Poxviridae. LSD, a major threat to stockbreeding, can cause acute or sub-acute disease in cattle and water buffalo (Givens 2018). All ages and breeds of cattle are affected; especially the young and cattle in the peak of lactation (Tuppurainen et al 2011). The reason why the World Organization for Animal Health (WOAH) has placed this transboundary disease on the notifiable disease list is due to its significant economic losses and the potential for rapid spread (Tuppurainen and Oura 2012). The recent spread of the disease in disease-free countries indicates the importance of its transmission as well as control and eradication (Sprygin et al 2019). It is transmitted by arthropod vectors such as mosquitoes, biting flies and ticks. In India, the disease was first

reported during the year 2019. The disease is characterized by mild fever for 2-3 days followed by development of stiff, round cutaneous nodules (2-5 cm in diameter) on the skin all over the body. These nodules are circumscribed, firm, round, raised and involve the skin, subcutaneous tissue and sometimes muscles. Symptoms may include lesions in mouth, pharynx and respiratory tract, emaciation, enlarged lymph nodes, oedema of limbs, reduction in milk production, abortion, infertility and sometimes, death. The morbidity rate is around 10-20 per cent and mortality rate is around 1-5 per cent (Arjkumpa et al 2022).

Causative agent: LSD is caused by the lumpy skin disease virus (LSDV) (Seerintra et al 2022), a member of the genus *Capripoxvirus* (CaPV) of the family Poxviridae. Capripoxviruses (CaPVs) represent one of the eight genera within the chordopoxvirus (ChPV) sub-family of the Poxviridae.

Lumpy skin disease and African swine fever spread across the country in less than two years after their first cases were reported



Source: African swine fever (ASF) situation update in Asia and Pacific, UN Food and Agriculture Organization, Media Reports

The *Capripoxvirus* genus currently comprises sheeppox virus (SPPV), goatpox virus (GTPV) and lumpy skin disease virus (LSDV), causing disease in sheep, goats or cattle respectively (Tulman et al 2002).

Transmission of lumpy skin disease: Long distance dispersal of LSDV seems to occur via the movement of infected animals, but distinct seasonal patterns indicate that arthropod-borne transmission is most likely responsible for the swift and aggressive short-distance spread of the disease. The mode of vector-borne transmission of the disease is most likely mechanical, but there is no clear-cut evidence to confirm or disprove this assumption. To date, the most likely vectors for LSDV transmission are blood-sucking arthropods such as stable flies (*Stomoxys calcitrans*), mosquitoes (*Aedes aegypti*) and hard ticks (*Rhipicephalus* and *Amblyomma* species). New evidence suggests that the ubiquitous,

synanthropic house fly, *Musca domestica* may also play a role in LSDV transmission, but this has not yet been tested in a clinical setting (Sprygin et al 2019). Infected bulls can excrete the virus in the semen, however, transmission of LSD via infected semen has not been demonstrated. It is not known if transmission can occur via fomites, for example ingestion of feed and water contaminated with infected saliva. Animals can be experimentally infected by being injected with blood or material from cutaneous nodules.

Clinical symptoms: Lumpy skin disease can occur in acute, sub-acute and chronic forms (Rouby et al 2021). The disease is characterized by mild fever for 2-3 days followed by development of stiff, round cutaneous nodules (2-5 cm in diameter) on the skin all over the body. These nodules are circumscribed, firm, round, raised and involve the skin, sub-cutaneous tissue and sometimes muscles (Hendrick 2017). The clinical

Endemic and Persistent

Diseases	Pathogens	Livestock	Zoonosis	Cases from 2017-2021
Anthrax	<i>Bacillus anthracis</i> bacteria	Multispecies	Yes	935
Babesiosis	<i>Babesia</i> parasite	Cattle	Yes*	16,185
Black Quarter	<i>Clostridium chauvoei</i> bacteria	Cattle and sheep	No	782
Bluetongue	Bluetongue virus	Ruminants, mainly sheep	No	3,349
Enterotoxaemia	<i>Clostridium perfringens</i> bacteria	Sheep and goat	No	2,012
Fasciolosis	<i>Fasciola hepatica</i> parasite	Cattle and sheep	Yes	15,292
Foot and mouth disease	Aphthovirus	Cattle, swine, sheep, goat	No	1,48,197
Hemorrhagic septicemia	<i>Pasteurella multocida</i> bacteria	Cattle and buffalo	No	5,122
Peste des petits ruminants	Morbillivirus	Goat and sheep	No	6,923
Sheep and Goat Pox	Sheep pox virus and goat pox virus	Sheep and goat	No	6,656
Swine fever	Pestivirus	Pigs	No	1,943
Theileriosis	<i>Theileria annulata</i> parasite	Cattle	No	2,719
Trypanosomosis	<i>Trypanosoma</i> parasite	Multispecies	Yes	6,556

INDIA

11.5% of income of semi medium and medium farmers (farmland 2.01-10 hectares) is from livestock farming

14.8% of income of small, marginal farmers (0.01-2 hectares) is from livestock farming

6.9% of income of large farmers (farmland over 10 hectares) is from livestock farming



WORLD

1.3 billion people depend on animals for their living

US \$300 billion is lost to livestock diseases each year

750 million of the world's poorest people depend on livestock support



Source: National Institute of Veterinary Epidemiology and Disease Informatics (NIVEDI): Land and Livestock Holdings of Households and Situation Assessment of Agricultural Households, 2019, Food and Agriculture Organization

features of the disease include fever, inappetence, nasal discharge, salivation and lacrymation, enlarged lymph nodes, a considerable reduction in milk production, loss of body weight and sometimes death (Tasioudi et al 2016). Some marks of crust are seen after bursting of the nodules on whole body. Symptoms may include lesions in mouth, pharynx and respiratory tract, emaciation, enlarged lymph nodes, edema in legs, dewlap and brisket region, reduction in milk production, abortion, infertility and sometimes death.

Postmortem findings: Due to lumpy skin disease, pox lesions can be seen all over the digestive and respiratory systems as well as on nearly any internal organs' surface. After the animal is skinned, sub-cutaneous lesions are easily discernible. There are

hemorrhages in the lungs, spleen and rumen. Distinctive skin nodules are also found. It includes ulcerative lesions in the respiratory and digestive tract mucosa; reddish, hemorrhagic to whitish lesions in the lungs, edoema (interlobular) and nodules in the lungs, heart lesions (endocardium) and skin vascular thrombosis followed by cutaneous infarction and sloughing (Singh 2022).

Diagnosis: Despite a primary clinical diagnosis of LSD, the diagnosis is confirmed by using conventional PCR (Zheng et al 2007) or real time PCR techniques (Bowden et al 2008). A real time PCR technique differentiates among LSDV, sheep and goat poxviruses (Lamien et al 2011). To differentiate virulent LSDV and vaccine strain, restriction fragment length

polymorphism (RFLP) has been used (Menasherow et al 2014). Stubbs et al (2012) stated that molecular methods are more precise, reliable and rapid compared to other methods. Among serological techniques, the virus neutralization test, though slow and costly with a high specificity and low sensitivity, is the only currently validated/valid test (Beard 2016).

Treatment: LSD being a viral illness, there is currently no specific treatment. LSD treatment is only symptomatic, with antibiotic medication which is used to prevent subsequent bacterial problems (Namazi and Tafti 2021). Antibiotics such as penicillins, cephalosporins, tetracyclines and fluroquinolones are prescribed for 5 to 7 days depending on the severity of the illness.

Salib and Osman (2011) conducted treatment studies for reducing LSD consequences. They were effective, utilizing a mix of medications that fight bacteria and inflammation, provide comfort and treat infections (Kumar et al 2022). They also recommended anti-histaminic and non-steroidal anti-inflammatory medications. An anti-pyretic medication, such as paracetamol, was given to reduce fever.

Abutarbush et al (2015) advised regular multivitamins and liver-supporting medications for anorexia recovery. Treating LSD and its effects is costly and does not always result in a full recovery; therefore, avoidance is more effective for minimizing large financial losses, milk loss from mastitis and losses of food products from death, miscarriage, fever and myiasis.

Economic impact: Cattle and buffalo play an important role in the socio-economic fabric of Asia. They are not only a source of income and food but also represent an accessible means to deposit savings and as a source of draught power. Such services help smallholders to overcome poverty hurdles, especially in poor settings where financial services and agricultural mechanization have not yet penetrated. Some communities value cattle and buffaloes highly for their role in social, cultural and religious traditions.

Knowing the economic impact of a disease can help decide on the most cost-effective disease control approach. Economically, Asia accounts for 31 and 98 per cent of global cow and buffalo milk production respectively and 29 per cent of the cattle

slaughtered globally for meat production (Roche et al 2020). LSD can inflict substantial direct losses through mortality, reduced milk production, damaged hides, poor growth, reduced draught power capacity and reproductive problems associated with abortions, infertility and lack of semen for artificial insemination. Vaccination costs, trade and other indirect revenue losses are directly proportional to the extent of LSD spread. LSD has led to serious economic losses in affected countries. The disease causes a considerable reduction in milk yield (from 10 to 85%) due to high fever and secondary mastitis (Namazi and Tafti 2021).

Other consequences of the disease include damaged hides, decline of the growth rate in beef cattle, temporary or permanent infertility, abortion, treatment and vaccination costs and death of infected animals (Sevik and Dogan 2017).

Limon et al (2020) in a study in northeast Nigeria, reported that lumpy skin disease (LSD), sheeppox (SP) and goatpox (GP) are contagious viral infections, affecting cattle (LSD), sheep and goats (SP and GP) with highly characteristic clinical signs affecting multiple body systems. All three diseases are widely reported to reduce meat, milk, wool and cashmere production although few studies have formally evaluated their economic impact on affected farms. The median incidence risk and fatality rate were 33 and 0 per cent in cattle, 53 and 34 per cent in sheep; 50 and 33 per cent in goats respectively, with young stock having higher incidence risk and fatality rates than adults. Almost all farmers (94%) treated affected animals with antibiotics, spending a median of USD 1.96 (min USD 0.19, max USD 27.5) per herd per day. Slaughtering or selling affected animals at low prices were common coping strategies. Milk production dropped 65 per cent when cows were clinically affected and 35 per cent after they recovered. Cattle lost a median of 10 per cent of their live weight and sheep and goats lost 15 per cent. Overall economic losses at farm level ranged from USD 9.6 to 6,340 depending on species affected and production system.

Chouhan et al (2022) determined the attack rate and risk factors along with economic losses of the LSD outbreaks in Bangladesh's Mymensingh and Gaibandha districts and reported that lumpy skin disease (LSD) is an emerging viral disease of cattle that negatively impacts livestock by reducing animals'

production and increasing management costs. The average economic loss per case was USD 110.40.

Kiplagat et al (2020), in a study in case-control study of cattle farms in Nakuru, Kenya, compared the economic impact between farms keeping purely indigenous or exotic breeds of cattle which indicated mean farm-level losses of USD 123 and 755 respectively. The mean farm-level losses from reduction in milk yield and mortality were estimated at USD 97 and 31 for farms keeping indigenous breeds whilst for farms keeping exotic breeds the equivalent losses were USD 266 and 431 respectively.

Klement (2018) reported that the incidence of LSD is the first factor which determines its direct economic impact. This depends on the abundance of vectors, the susceptibility of the host and the use of preventive measures (Gari et al 2011). It can reach even 85 per cent in an affected herd if no preventive measures are applied (Tuppurainen and Oura 2012). Case fatality is also an important factor, influencing the economic impact of a disease. However, accurate estimation of case fatality is very difficult to provide as in most of the developed countries, sick animals are culled and in developing countries, the exact pathological reason for natural animal death is not always provided.

Prevention and control: The treatment of LSD is only symptomatic and targeted at preventing secondary bacterial complications using a combination of antimicrobials, anti-inflammatory, supportive therapy and antiseptic solutions (Salib and Osman 2011). Vaccination is the only effective method to control the disease in endemic areas along with movement restrictions and the removal of affected animals (Sevik and Dogan 2017). The culling of affected animals, movement restrictions and compulsory and consistent vaccination have been recommended as control strategies (Beard 2016). Moreover, risk factors should be considered in control activities (Sevik and Dogan 2017).

Educating veterinarians and livestock workers would enable them to perform timely diagnoses of clinical cases, helping to slow the spread of disease (Beard 2016). However, regarding the role of arthropod vectors, elimination of the disease is likely to be difficult and any delays in the removal of infected animals increase the risk of LSD transmission (Tuppurainen et al 2017).

Furthermore, the rapid confirmation of a clinical diagnosis is essential so that eradication measures, such as quarantine, slaughter out of affected and in contact animals, proper disposal of carcasses, cleaning and disinfection of the premises and insect control can be implemented as soon as possible during the eruption (Constable et al 2016). Moreover, rigorous import restrictions on livestock, carcasses, hides and semen from endemic areas must be in place in disease free areas (Sevik and Dogan 2017).

Vaccination of cattle using a vaccine with demonstrated efficacy is the best option for controlling the spread of LSD, especially if pre-emptive ie applied before the virus enters a region or country at risk. However, preventive vaccination against LSD leads to trade restrictions on the export of live cattle and their products, which may deter disease-free exporting countries from implementing pre-emptive vaccination in high-risk regions. Pre-emptive vaccination is highly recommended when LSD is detected across borders in neighbouring countries. It may take the form of zone or buffer vaccination, taking into account geographical barriers, transport access routes and host population densities (Anon 2017). The commercially accessible vaccines against LSD are live attenuated vaccines.

Vaccine and vaccination: Vaccination is the only effective method of control the disease in the endemic area along with movement restrictions and removal of affected animals. Bazid et al (2022) conducted a study to evaluate the safety and efficacy of a new live attenuated LSD vaccine produced by Middle East for vaccines (MEVAC®) based on the Neethling strain. In the field, around 2 per cent (80/4301) of the animals showed hyper-reactivity and 0.6 per cent (24/4301) showed small skin swellings that disappeared within few hours PV. Abortion was recorded in three animals (0.3%, 3/867).

In LSD free countries that use the sheep pox vaccine to protect sheep against sheep pox, it was recommended to use the same vaccine during LSD outbreaks because of potential safety issues associated with the live attenuated LSDV vaccine use (Tuppurainen and Oura 2012). Live vaccines produce a strong and long lasting immune response and are efficient in the control of disease spread (Tuppurainen et al 2020). However, live vaccines can cause local inflammation and a mild disease with skin lesions (Bedekovic et al 2017). Inactivated vaccines could be applied in the final stage of disease eradication as a

part of the strategy that uses live vaccines first (Hamdi et al 2020).

As there is a chance of recombination between the wild field strain and the live vaccine, the risk of coinfection should be considered with the use of live vaccines and natural infection is probably made worse by the vaccination of infected animals (Sprygin et al 2019).

Vaccination of cattle using a vaccine with demonstrated efficacy is the best option for controlling the spread of LSD, especially if pre-emptive is applied before the virus enters a region or country at risk. However, preventive vaccination against LSD leads to trade restrictions on the export of live cattle and their products, which may deter disease-free exporting countries from implementing pre-emptive vaccination in high-risk regions. Pre-emptive vaccination is highly recommended when LSD is detected across borders in neighbouring countries. It may take the form of zone or buffer vaccination, taking into account geographical barriers, transport access routes and host population densities (Anon 2017).

The ten-fold dose of attenuated SPPV vaccines is recommended for immunization of bovines against LSD. Nevertheless, compared to the Neethling vaccine its efficacy is significantly lower (Ben-Gera et al 2015). Currently, most commercially available vaccines against LSD are live attenuated vaccines based on a LSDV strain, sheeppox virus (SPPV), or goatpox virus (GTPV) (Tuppurainen et al 2021). Gorgan GTPV vaccine is a good, cost-effective alternative in those countries where GTP and LSD overlap. Recent data from Kazakhstan also enhance the understanding of the potential use of GTPV where the goatpox vaccine strain for LSD elicited a strong protective immune response in cattle (Zhugunissov et al 2020). In India the GTPV Uttarkashi strain is being evaluated for level of protection against LSD as compared to the LSDV vaccine and is already used for emergency vaccination. In Bangladesh, GTPV vaccine was used in Chattogram and found to be effective against LSD (Kayesh et al 2020). There are also several studies of the GTPV vaccine based on the Gorgan strain with successful results (Zhugunissov et al 2020).

Self-observed future strategies to control lumpy disease in livestock: Controlling lumpy disease in animals requires a comprehensive approach that

includes preventive measures, early detection and appropriate management strategies. As of last knowledge update in September 2021, there was no specific information available on a disease called lumpy disease in animals. However, assuming to a hypothetical condition or a specific disease that may emerge in the future, here are some general strategies that can be applied to control and manage infectious diseases in animals:

- Vaccination programmes
- Quarantine and biosecurity
- Surveillance and early detection
- Rapid response and containment
- Education and awareness
- Research and development
- Collaboration and coordination

It is important to note that specific strategies for controlling a particular disease may vary depending on its characteristics, including the mode of transmission, affected animal species, geographical location and available resources. Therefore, it is crucial to consult veterinary experts and authorities to tailor the control measures to the specific disease.

CONCLUSION

The recent spread of the disease into disease free areas indicates its epidemiological and economic significance. Considering the extensive boundaries of Middle East countries, animal movements among these countries should be attentively controlled by veterinary authorities. Furthermore, paying close attention to the different aspects of the disease, such as transmission and epidemiology and the implementation of effective preventive measures such as vaccination, could result in better disease control. Therefore, accurate and timely diagnosis in endemic areas, vaccination with the homologous strain of the LSDV, vector control, animal movement restriction and LSDV testing of bulls used for breeding are highly recommended as tools to control further spread.

REFERENCES

- Abutarbush SM, Ababneh MM, Al Zoubi IG, Al Sheyab OM, Al Zoubi MG, Alekish MO and Al Gharabat RJ 2015. Lumpy skin disease in Jordan: disease emergence, clinical signs, complications and preliminary associated economic losses. *Transboundary and Emerging Diseases* **62(5)**: 549-554.

- Anonymous 2017. Sustainable prevention, control and elimination of lumpy skin disease – eastern Europe and the Balkans. FAO Animal Production and Health Position Paper Number 2, Rome, Italy.
- Arjkumpa O, Suwannaboon M, Boonrod M, Punyawan I, Liangchaisiri S, Laobannue P, Lapchareonwong C, Sansri C, Kuatako N, Panyasomboonying P, Uttarat P, Buamithup N, Sansamur C and Punyapornwithaya V 2022. The first lumpy skin disease outbreak in Thailand (2021): epidemiological features and spatio-temporal analysis. *Frontiers in Veterinary Science* **8**: 799065; doi: 10.3389/fvets.2021.799065.
- Bazid A-H, Wasfy M, Fawzy M, Nayel M, Abdelmegeid M, Thabet RY, Yong HS, El-Sayed MM, Magouz A and Badr Y 2022. Emergency vaccination of cattle against lumpy skin disease: evaluation of safety, efficacy and potency of MEVAC® LSD vaccine containing Neethling strain. *Veterinary Research Communications*, doi:10.1007/s11259-022-10037-2.
- Beard PM 2016. Lumpy skin disease: a direct threat to Europe. *Veterinary Record* **178**(22): 557-558.
- Bedekovic T, Simic I, Kresic N and Lojkic I 2017. Detection of lumpy skin disease virus in skin lesions, blood, nasal swabs and milk following preventive vaccination. *Transboundary and Emerging Diseases* **65**(2): 491-496.
- Ben-Gera J, Klement E, Khinich E, Stram Y and Shpigel NY 2015. Comparison of the efficacy of Neethling lumpy skin disease virus and x10RM65 sheep-pox live attenuated vaccines for the prevention of lumpy skin disease – the results of a randomized controlled field study. *Vaccine* **33**(38): 4837-4842.
- Bowden TR, Babiuk SL, Parkyn GR, Copps JS and Boyle DB 2008. Capripoxvirus tissue tropism and shedding: a quantitative study in experimentally infected sheep and goats. *Virology* **371**(2): 380-393.
- Chouhan CS, Parvin MS, Ali MY, Sadekuzzaman M, Chowdhury MGA, Ehsan MA and Islam MT 2022. Epidemiology and economic impact of lumpy skin disease of cattle in Mymensingh and Gaibandha districts of Bangladesh. *Transboundary and Emerging Diseases* **69**(6): 3405-3418.
- Constable PD, Hinchcliff KW, Done SH and Grundberg W 2016. *Veterinary medicine: a textbook of the diseases of cattle, horses, sheep, pigs and goats*. 11th Edn, Elsevier, 2278p.
- Gari G, Bonnet P, Roger F, Waret-Szkuta A 2011. Epidemiological aspects and financial impact of lumpy skin disease in Ethiopia. *Preventive Veterinary Medicine* **102**: 274-283.
- Givens MD 2018. Review: risks of disease transmission through semen in cattle. *Animals* **12**(s1): s165-s171.
- Hamdi J, Boumart Z, Daouam S, El Arkam A, Bamouh Z, Jazouli M, Tadlaoui KO, Fihri OF, Gavrilov B and El Harrak M 2020. Development and evaluation of an inactivated lumpy skin disease vaccine for cattle. *Veterinary Microbiology* **245**: 108689; doi: 10.1016/j.vetmic.2020.108689.
- Hendrick MJ 2017. Mesenchymal tumors of the skin and soft tissues. In: *Tumors in domestic animals*, 5th Edn, (DJ Meuten, Ed), John Wiley and Sons Inc, Ames, Iowa, USA, pp 142-175.
- Kayesh MEH, Hussan MT, Hashem MA, Eliyas M and Anower AKMM 2020. Lumpy skin disease virus infection: an emerging threat to cattle health in Bangladesh. *Hosts and Viruses* **7**(4): 97-108.
- Kiplagat SK, Kitale PM, Onono JO, Beard PM and Lyons NA 2020. Risk factors for outbreaks of lumpy skin disease and the economic impact in cattle farms of Nakuru County, Kenya. *Frontiers in Veterinary Science* **7**: 259; doi: 10.3389/fvets.2020.00259.
- Klement E 2018. Economic impact of lumpy skin disease. In: *Lumpy skin disease* (ESM Tuppurainen, S Babiuk and E Klement, Eds), Springer, Cham, pp 7-9.
- Kumar N, Barua S, Kumar R, Khandelwal N, Kumar A, Verma A, Singh L, Godara B, Chander Y, Kumar G, Riyesh T, Sharma DK, Pathak A, Kumar S, Dedar RK, Mehta V, Gaur M, Bhardwaj B, Vyas V, Chaudhary S, Yadav V, Bhati A, Kaul R, Bashir A, Andrabi A, Yousuf RW, Koul A, Kachhawaha S, Gurav A, Gautam S, Tiwari HA, Munjal VK, Gupta MK, Kumar R, Gulati BR, Misri J, Kumar A, Mohanty AK, Nandi S, Singh KP, Pal Y, Dutt T and Tripathi BN 2022. Evaluation of the safety, immunogenicity and efficacy of a new live-attenuated lumpy skin disease vaccine in India. *Virulence* **14**(1): 2190647; doi: 10.1080/21505594.2023.2190647.
- Lamien CE, Lelemta M, Goger W, Silber R, Tuppurainen E, Matijevic M, Luckins AG and Diallo A 2011. Real time PCR method for simultaneous detection, quantitation and differentiation of capripoxviruses. *Journal of Virological Methods* **171**(1): 134-140.
- Limon G, Gamawa AA, Ahmed AI, Lyons NA and Beard PM 2020. Epidemiological characteristics and economic impact of lumpy skin disease, sheeppox and goatpox among subsistence farmers in northeast Nigeria. *Frontiers in Veterinary Science* **7**: 8; doi: 10.3389/fvets.2020.00008.
- Menasherow S, Rubinstein-Giuni M, Kovtunen A, Eynor Y, Fridgut O, Rotenberg D, Khinich Y and Stram Y 2014. Development of an assay to differentiate between

- virulent and vaccine strains of lumpy skin disease virus (LSDV). *Journal of Virological Methods* **199**: 95-101.
- Namazi F and Tafti AK 2021. Lumpy skin disease, an emerging transboundary viral disease: a review. *Veterinary Medicine and Science* **7(3)**: 888-896.
- Roche X, Rozstalnyy A, Tago Pacheco D, Kamata A, Pittiglio C, Alcrudo DB, Bisht K, Karki S, Kayamori J, Larfaoui F, Raizman E, VonDobschuetz S, Dhingra MS and Sumption K 2020. Introduction and spread of lumpy skin disease in south, east and southeast Asia: qualitative risk assessment and management. *FAO Animal Production and Health Paper 183*, Food and Agriculture Organization of the United Nations, Rome, Italy, 62p.
- Rouby SR, Shehata O, Abdel-Moneim AS, Hussein KH and Mahmoud MM 2021. Lumpy skin disease in calves: the association between clinical signs and biochemical alterations. *Advances in Animal and Veterinary Sciences* **9(11)**: 1863-1868.
- Salib FA and Osman AH 2011. Incidence of lumpy skin disease among Egyptian cattle in Giza Governorate, Egypt. *Veterinary World* **4(4)**: 162-167.
- Seerintra T, Saraphol B, Wankaew S and Piratae S 2022. Molecular identification and characterization of lumpy skin disease virus emergence from cattle in the northeastern part of Thailand. *Journal of Veterinary Science* **23(5)**: e73; doi: 10.4142/jvs.22111.
- Sevik M and Dogan M 2017. Epidemiological and molecular studies on lumpy skin disease outbreaks in Turkey during 2014-2015. *Transboundary and Emerging Diseases* **64(4)**: 1268-1279.
- Singh R 2022. Lumpy skin disease in bovines: a dreadful emerging disease in cattle. <https://www.pashudhanpraharee.com/lumpy-skin-disease-in-bovines/> (Retrieved: 12.04.2023)
- Sprygin A, Pestova Y, Wallace DB, Tuppurainen E and Kononov AV 2019. Transmission of lumpy skin disease virus: a short review. *Virus Research* **269**: 197637; doi: 10.1016/j.virusres.2019.05.015.
- Stubbs S, Oura CA, Henstock M, Bowden TR, King DP and Tuppurainen ES 2012. Validation of a high throughput real time polymerase chain reaction assay for the detection of capripoxviral DNA. *Journal of Virological Methods* **179(2)**: 419-422.
- Tasioudi KE, Antoniou SE, Iliadou P, Sachpatzidis A, Plevraki E, Agianniotaki EI, Fouki C, Mangana-Vougiouka O, Chondrokouki E and Dile C 2016. Emergence of lumpy skin disease in Greece, 2015. *Transboundary and Emerging Diseases* **63(3)**: 260-265.
- Tulman ER, Afonso CL, Lu Z, Zsak L, Sur J-H, Sandybaev NT, Kerembekova UZ, Zaitsev VL, Kutish GF and Rock DL 2002. The genomes of sheeppox and goatpox viruses. *Journal of Virology* **76(12)**: 6054-6061.
- Tuppurainen ESM and Oura CAL 2012. Review: lumpy skin disease: an emerging threat to Europe, the Middle East and Asia. *Transboundary and Emerging Diseases* **59(1)**: 40-48.
- Tuppurainen ESM, Antoniou SE, Tsiamadis E, Topkaridou M, Labus T, Debeljak Z, Plavsic B, Miteva A, Alexandrov T, Pite L, Boci J, Marojevic D, Kondratenko V, Atanasov Z, Murati B, Acinger Rogic Z, Kohnle L, Calistri P and Broglia A 2020. Field observations and experiences gained from the implementation of control measures against lumpy skin disease in South East Europe between 2015 and 2017. *Preventive Veterinary Medicine* **181**: 104600; doi: 10.1016/j.prevetmed.2018.12.006.
- Tuppurainen ESM, Dietze K, Wolff J, Bergmann H, Beltran-Alcrudo D, Fahrion A, Lamien CE, Busch F, Sauter-Louis C, Conraths FJ, De Clercq K, Hoffman B and Knauf S 2021. Review: vaccines and vaccination against lumpy skin disease. *Vaccines* **9(10)**: 1136; doi: 10.3390/vaccines9101136.
- Tuppurainen ESM, Stoltz WH, Troskie M, Wallace DB, Oura CAL, Mellor PS, Coetzer JAW and Venter EH 2011. A potential role for ixodid (hard) tick vectors in the transmission of lumpy skin disease virus in cattle. *Transboundary and Emerging Diseases* **58(2)**: 93-104.
- Tuppurainen ESM, Venter EH and Coetzer JAW 2017. The detection of lumpy skin disease virus in samples of experimentally infected cattle using different diagnostic techniques. *Onderstepoort Journal of Veterinary Research* **72(2)**: 153-164.
- Zheng M, Liu Q, Jin NY, Guo JG, Huang X, Li HM, Zhu W and Xiong Y 2007. A duplex PCR assay for simultaneous detection and differentiation of *Capripoxvirus* and orf virus. *Molecular and Cellular Probes* **21(4)**: 276-281.
- Zhugunisssov K, Bulatov Y, Orynbayev M, Kutumbetov L, Abduraimov Y, Shaiyakhmetov Y, Taranov D, Amanova Z, Mambetaliyev M, Abstova Z, Azanbekova M, Khairullin B, Zakarya K and Tuppuainen E 2020. Goatpox virus (G20-LKV) vaccine strain elicits a protective response in cattle against lumpy skin disease at challenge with lumpy skin disease virulent field strain in a comparative study. *Veterinary Microbiology* **245**: 108695; doi: 10.1016/j.vetmic.2020.108695.