

Wild edible fruit plants used by rural people of outer Seraj region of district Kullu, Himachal Pradesh

NISHA*, RAJ KUMAR and RHYTHM GOSWAMI

ICFRE – Himalayan Forest Research Institute, Conifer Campus
Panthaghati, Shimla 171013 Himachal Pradesh, India

*Email for correspondence: presstonisha@gmail.com/nisha2092sharma@gmail.com

© Society for Advancement of Human and Nature (SADHNA)

Received: 19.02.2024/Accepted: 13.04.2024

ABSTRACT

Wild edible fruits can play a significant role in providing nutrition to the rural inhabitants. The present study was carried out in outer Seraj region of district Kullu, Himachal Pradesh. The data were collected through interviews, group discussions, participatory observations and semi-structured questionnaire. This survey documented total of 24 wild edible fruit plants which belonged to different categories viz 16 trees, 7 shrubs and 1 herb. These documented plants were used by the rural people of the study area as wild fruits and vegetables. It was found that the highest number (6) of wild edible plants belonged to Rosaceae family followed by Polygonaceae (4) and Berberidaceae (2) families. These wild plants were easily available from the surrounding without any cost. These plants have been reported to be of high nutritional values. Due to unscientific and ruthless exploitation coupled with habitat degradation, these species have been declining very fast. Biodiversity of all these species needs to be conserved by adopting different management practices so as to save them from extinction in the near future.

Keywords: Wild fruits; rural people; habitat degradation; biodiversity

INTRODUCTION

Wild edible plants have played a significant role in meeting the nutritional requirement and livelihood improvement of the rural inhabitants since time immemorial. These plants have been utilized for food, fibre, medicine and fodder for domestic animals (Janjua et al 2020). The utilization of wild edible plants as a nutritional supplement has a lot of potential (Prakash et al 2021). Therefore, these plants can increase quality of daily food for the rural population. The Himalayas are popularly known for their abundance of wild edible plants. The edible plants also contribute significantly for supporting the nutritional requirements of the indigenous population in distant areas of the country throughout the year (Thakur 2017). Hence, wild edible fruit plants can be potential food supplements so as to increase quality of daily food for the rural population.

In India, various studies have been carried out on the utilization pattern of wild edible plants (Satyavani et al 2015, Rashid et al 2008, Sankaran et al 2006,

Upadhye et al 1994, Radha et al 2021, Valvi et al 2011, Thakur 2017).

In Himachal Pradesh, the documentation of wild edible plant diversity has been carried out through a number of studies (Jain 1991, Maikhuri et al 1998, Samant et al 1998, Chauhan 1999, Joshi et al 1999, Sood et al 2001, Dhar et al 2002, Samant and Pal 2003, Seth and Jaswal 2004, Rawat and Garg 2005, Kala 2006, Samant and Pant 2006, Uniyal et al 2006, Samant et al 2007, Rawat et al 2009, Singh et al 2009, Semwal et al 2010).

Outer Seraj area, having diverse climatic and topographic conditions, provides a diverse range of conditions for the luxuriant growth of potential medicinal plants (Singh and Rawat 1999). Indigenous people of this region collect wild edible plants for food and other plants from natural areas to fulfil their subsistence needs, in addition to planting a few crops. These fruit plants are threatened by population pressure and man's activities like grazing and burning. In addition, due to

westernization, new generation is not much interested in traditional knowledge of plants so there is urgent need to document traditional knowledge of wild edible plants from this unexplored region of Himachal Pradesh. This region also requires further exploration in order to encourage the domestication of commercially relevant species. Therefore, there is an urgent need to document various important wild edible plant species of this area. This study would be helpful for future researchers and scientists working in the field of systematic, biochemical and pharmacological studies.

METHODOLOGY

Outer Seraj is one of the most beautiful regions of Kullu district blessed with beautiful flora and fauna. It is situated between 31°39'32" N latitude and 77°23'12" E longitudes and elevation range from 700 m to 5,500 m amsl (Fig 1). The study area comprised 60 villages of Anni and Nirmand blocks of Kullu district.

The present study included extensive field surveys conducted in the year 2022 and 2023. In current study, villagers were surveyed at household

level and data were collected through interviews, group discussions, participatory observations and semi-structured questionnaire. Information was obtained on local names, part(s) used and indigenous uses of the plants. The informants included men, women, youths and elders between the age of 28 and 75 years. The information was documented and analyzed for various parameters. Collection of fresh samples was done and samples were identified with the help of sources on local flora (Collett 1902, Aswal and Mehrotra 1994, Dhaliwal and Sharma 1999, Singh and Rawat 2000).

RESULTS and DISCUSSION

In current the study a total of 24 edible wild fruit plants were reported from the outer Seraj area (Table1). The detailed information regarding local name, medicinal use of plant species and their mode of application is presented. The documented 24 plant species from the study area belonged to 13 families and 18 genera. It was observed that most of the plant species were reported from Rosaceae, Moraceae and Berberidaceae families. Rosaceae family was represented by total 6 species followed by Moraceae by 4, Berberidaceae by 2, while the other families




Fig 1. Map of outer Seraj region depicting study area

Table 1. Wild edible fruit plants and their uses in outer Seraj, Kullu, Himachal Pradesh

Botanical name	Family	Local name	Habit	Edible part(s)	Mode of consumption	Medicinal use(s)
<i>Aegle marmelos</i> (L) Corrêa	Rutaceae	Bel	Tree	Fruit pulp	Natural drink prepared from mucilaginous pulp	Treating dysentery, vomiting, fever
<i>Aesculus indica</i> (Wall ex Cambess) Hook	Sapindaceae	Khanor	Tree	Ripe fruits	Dried fruits crushed into flour and used as tonic	Cures body weakness, menstrual problems
<i>Berberis aristata</i> DC	Berberidaceae	Kashmal	Shrub	Ripe fruits	Fruits edible	Jaundice, piles
<i>Berberis lycium</i> Royle	Berberidaceae	Kashmal	Shrub	Ripe fruits	Fruits eaten	Jaundice, cough, malaria, cold
<i>Carissa spinarum</i> L	Apocynaceae	Karonda	Shrub	Ripe fruits	Ripe fruits eaten	Headache, rheumatism
<i>Cordia dichotoma</i> G Forst	Cordiaceae	Lasura	Tree	Ripe fruits	Mucilage from ripe fruits used as gum	Cough, cold
<i>Ficus auriculata</i> Lour	Moraceae	Trimbal	Tree	Ripe, unripe fruits	Unripe fruits used as vegetable; ripe fruits eaten	Dysentery, diabetes, piles
<i>Ficus carica</i> L	Moraceae	Fedu	Tree	Ripe, unripe fruits	Unripe fruits used as vegetable; ripe fruits eaten	Diabetes
<i>Ficus palmata</i> Forsk	Moraceae	Fegda	Tree	Ripe, unripe fruits	Unripe fruits used as vegetable; ripe fruits eaten	Constipation
<i>Ficus racemosa</i> L.	Moraceae	Gular	Tree	Ripe, unripe fruits	Unripe fruits used as vegetable; ripe fruits eaten	Diabetes, liver disorders
<i>Flacourtia indica</i> (Burm f) Merr	Salicaceae	Kangru	Tree	Ripe fruits	Acidic ripe fruits eaten	Treating diarrhoea
<i>Fragaria indica</i> Wall	Rosaceae	Bhumlae	Herb	Ripe fruits	Ripe fruits edible	Blood purifier
<i>Juglans regia</i> L	Juglandaceae	Khod	Tree	Ripe fruits	Nuts consumed after sun drying	Skin disorders, arthritis
<i>Myrica esculenta</i> Buch-Ham ex D Don	Myricaceae	Kaphal	Tree	Ripe fruits	Ripe fruits consumed directly	Asthma, cough, inflammation
<i>Phyllanthus emblica</i> L	Phyllanthaceae	Aonla	Tree	Ripe fruits	Ripe fruits edible	Stomach-ache, eye complaints, fever
<i>Princepia utilis</i> Royle	Rosaceae	Bhekhal	Shrub	Ripe fruits	Ripe fruits edible	Cough, cold, high blood pressure

Botanical name	Family	Local name	Habit	Edible part(s)	Mode of consumption	Medicinal use(s)
<i>Prunus armeniaca</i> L	Rosaceae	Khumani	Tree	Ripe fruits	Ripe fruits sweet in taste, eaten	To treat constipation
<i>Prunus cornuta</i> (Wall ex Royle) Steud	Rosaceae	Himalayan bird cherry	Tree	Ripe fruits	Ripe fruits eaten	Digestive problems, skin diseases
<i>Punica granatum</i> L	Lythraceae	Daru, Anar	Tree	Ripe fruits	Ripe fruits edible	Diabetes
<i>Pyrus pashia</i> Buch-Ham ex D Don	Rosaceae	Kainth	Tree	Ripe fruits	Ripe fruits eaten	Conjunctivitis
<i>Rubus ellipticus</i> Sm	Rosaceae	Hinsar	Shrub	Ripe fruits	Ripe fruits eaten	Bronchitis and diabetes
<i>Rubus niveus</i> Thunb	Rosaceae	Aakhe	Shrub	Ripe fruits	Ripe fruits eaten	Antistress, antibacterial
<i>Syzygium cumini</i> (L) Skeels	Myrtaceae	Jamun	Tree	Ripe fruits	Ripe fruits eaten	Diabetes
<i>Ziziphus nummularia</i> (Burm f) Wight & Arn	Rhamnaceae	Ber	Shrub	Ripe fruits	Ripe fruits sweet in taste, edible	Indigestion, dysentery

reported during the current survey represented 1 species each (Fig 2).

It was observed that most of the edible plant species reported from outer Seraj region were trees (66.67%) followed by shrubs (29.17%) and herb (4.2%) (Fig 3). The ripe fruits were most commonly used as edible plant part followed by fruit pulp. These wild fruits are of great importance due to their medicinal properties. The common diseases people used to treat were cold, cough, fever, constipation, wounds, joint pain, dysentery, jaundice, piles, diabetes etc. Some important edible wild fruit plants of outer Seraj area are shown in Plate 1.

Wild plants have shown an important role among rural inhabitants that relied on wild food resources for food and medicinal purposes. The people of this area had been living in isolated, mountainous region or close to forest areas, where no proper transportation and healthcare facilities were accessible. Research on wild edible fruit species was intended to promote the preservation of these twenty four species and for welfare of local communities in order to supply their nutritional requirements. In addition to their nutritional value, the preservation of these underutilized edible fruits also has economical advantages.

An attempt can be made to utilize these edible wild fruits as a source of income, particularly for poor

rural poor and unemployed youth. With little efforts, a variety of products can be made from the wild edible fruits and supplied to food processing industries. The medicinal values of these fruits as tonics and appetizers have further scope of their popularity. There is a need to conduct systematic, biochemical and pharmacological studies to validate the traditional claims.

CONCLUSION

From the present study, it is evident that underutilized wild edible fruit plants provide nutrition to the forest dwellers and many of the marginalized rural inhabitants since the commonly cultivable fruits are costly and unaffordable for them. However, due to modernization, there has been continuous decline in the use of locally available resources as a source of food and relying entirely on staple food plants. There is an urgent need to investigate how to reduce the current reliance on staple crops to meet the needs of the society. During the study, it was found that most of the fruits documented were normally eaten raw when they were ripe, whereas, unripe fruits were usually cooked as vegetables. It is recommended that documentation and conservation of wild edible plants is very important before these are eliminated from the society. It is feared that they will become extinct in due course of time. This study would also be helpful for food industry to develop new edible food products.




Fig 2. Wild edible fruit plants of outer Seraj, Kullu as per families




Fig 3. Habit-wise distribution of wild edible fruit plants of outer Seraj, Kullu

ACKNOWLEDGEMENT

Authors express their gratitude to the rural people of outer Seraj region of Kullu district, Himachal Pradesh for sharing their empirical knowledge about wild edible fruit plants of the study area.

REFERENCES

- Aswal BS and Mehrotra BN 1994. Flora of Lahaul-Spiti – a cold desert in northwest Himalaya. Bishen Singh Mahendra Pal Singh, Dehradun, Uttarakhand, India, 761p.
- Chauhan NS 1999. Medicinal and aromatic plants of Himachal Pradesh. Indus Publishing Company, New Delhi, India, 632p.
- Collett H 1902. Flora simlensis – a handbook of the flowering plants of Simla and the neighbourhood. Thacker Spink and Company, London.
- Dhaliwal DS and Sharma M 1999. Flora of Kullu district (Himachal Pradesh). 1st Edn, Bishen Singh Mahendra Pal Singh, Dehradun, Uttarakhand, India.
- Dhar U, Manjkhola S, Joshi M, Bhatt A, Bisht AK and Joshi M 2002. Current status and future strategy for


*Fragaria indica**Pyrus pashia**Ficus racemosa**Berberis aristata**Myrica esculenta**Prunus cornuta*

Plate 1. Some important wild edible fruit plants of outer Seraj region of Kullu district, Himachal Pradesh

development of medicinal plants sector in Uttaranchal, India. *Current Science* **83**(8): 956-964.

Jain SK 1991. Dictionary of Indian folk medicine and ethnobotany – a reference manual of man-plant relationships, ethnic groups and ethnobotanists in India. Deep Publications, New Delhi, India, 311p.

Janjua S, Srivastava S and Negi V 2020. Ethnobotanical study of medicinal plants used in Shikari Devi Wildlife Sanctuary of Himachal Pradesh, India. *Medicinal Plants – International Journal of Phytomedicines and Related Industries* **12**(4): 666-673.

Joshi HC, Arya SC and Samant SS 1999. Diversity, distribution and indigenous uses of medicinal and edible plants in a part of Nanda Devi Biosphere Reserve – I. *Himalayan Biosphere Reserves* **1**(1-2): 49-65.

Kala CP 2006. Medicinal plants of the high altitude cold desert in India: diversity, distribution and traditional uses. *International Journal of Biodiversity Science and Management* **2**(1): 43-56.

Maikhuri RK, Nautiyal S, Rao KS and Saxena KG 1998. Role of medicinal plants in the traditional healthcare system: a case study from Nanda Devi Biosphere Reserve. *Current Science* **75**(2): 152-157.

Prakash P, Radha, Kumar M, Kumari N, Prakash S, Rathour S, Thakur M, Jamwal R, Janjua S, Ali M, Pundir A, Puri S, Dhumal S, Singh S, Senapathy M, Bangar SP, Maurya VK, Changan S, Gora JS, Samota MK, Damale RD, Sasi M, Natta S, Chandran D, Rajalingam S, Rais N and Mekhemar M 2021. Therapeutic uses of wild plants by rural inhabitants of Maraog region in district Shimla, Himachal Pradesh, India. *Horticulturæ* **7**(10): 343; doi: 10.3390/horticulturæ7100343.

Radha, Janjua S, Ali M, Thakur M, Jamwal R, Rathour S, Pubral AK, Kumari N, Puri S, Pundir A and Kumar M 2021. Documenting traditional knowledge before they are forgotten: a study on the ethnomedicinal uses of wild plants by rural people of Jubbarhatti in district Shimla, Himachal Pradesh, India. *International Journal of Theoretical and Applied Sciences* **13**(1): 37-51.

Rashid A, Anand VK and Serwar J 2008. Less known wild edible plants used by the Gujjar tribe of district Rajouri, Jammu and Kashmir state. *International Journal of Botany* **4**(2): 219-224.

Rawat RBS and Garg GP 2005. Medicinal plants: trade and commerce opportunities with India. *Indian Forester* **131**(3): 275-287.

Rawat RBS, Jishtu V and Kapoor KS 2009. Medicinal and aromatic plant diversity of Himalayan cold desert with

- reference to Spiti valley of northwest Himalayas. *Indian Forester* **135(7)**: 891-904.
- Samant SS and Pal M 2003. Diversity and conservation status of medicinal plants in Uttaranchal state. *Indian Forester* **129(9)**: 1090-1108.
- Samant SS and Pant S 2006. Diversity, distribution pattern and conservation status of the plants used in liver diseases/ailments in Indian Himalayan region. *Journal of Mountain Science* **3(1)**: 28-47.
- Samant SS, Dhar U and Palni LMS 1998. Medicinal plants of Indian Himalaya: diversity distribution potential values. GB Pant Institute of Himalayan Environment and Development, Almora, Uttarakhand, India, 163p.
- Samant SS, Pant S, Singh M, Lal M, Singh A, Sharma A and Bhandari S 2007. Medicinal plants in Himachal Pradesh, northwestern Himalaya, India. *International Journal of Biodiversity Science and Management* **3(4)**: 234-251.
- Sankaran M, Prakash J, Singh NP and Suklabaidya A 2006. Wild edible fruits of Tripura. *Natural Product Radiance* **5(4)**: 302-305.
- Satyavani S, Satyavathi K, Devisoundarya S and Padal SB 2015. Wild edible plants used for indigenous community of north-coastal Andhra Pradesh, India. *BMR Phytomedicine* **2(1)**: 1-7.
- Semwal DP, Saradhi PP, Kala CP and Sajwan BS 2010. Medicinal plants used by local Vaidyas in Ukhimath block, Uttarakhand. *Indian Journal of Traditional Knowledge* **9(3)**: 480-485.
- Seth MK and Jaswal S 2004. An enumeration of plant resources of Shimla (Himachal Pradesh) in the northwest Himalayas. International Book Distributors and Publishers, Dehradun, Uttarakhand, India.
- Singh A, Lal M and Samant SS 2009. Diversity, indigenous uses and conservation prioritization of medicinal plants in Lahaul valley, proposed Cold Desert Biosphere Reserve, India. *International Journal of Biodiversity Science and Management* **5(3)**: 132-154.
- Singh SK and Rawat GS 1999. Floral diversity and vegetation structure in Great Himalayan National Park, western Himalaya. FREEP-GHNP 02/05, FREEP-GHNP Research Project, Wildlife Institute of India, Dehradun, Uttarakhand, India, 103p.
- Singh SK and Rawat GS 2000. Flora of Great Himalayan National Park, Himachal Pradesh. 1st Edn, Bishen Singh Mahendra Pal Singh, Dehradun, Uttarakhand, India.
- Sood SK, Nath R and Kalia DC 2001. Ethnobotany of cold desert tribes of Lahaul-Spiti (NW Himalaya). Deep Publications, New Delhi, India.
- Thakur SD 2017. Diversity, distribution and utilization pattern of some forestry foods (wild edibles) from Tirthan Wildlife Sanctuary of dist Kullu, Himachal Pradesh. *International Journal of Advances in Science Engineering and Technology* **5(2) (Special Issue 2)**: 4-11.
- Uniyal SK, Kumar A, Lal B and Singh RD 2006. Quantitative assessment and traditional uses of high value medicinal plants in Chhota Bhangal area of Himachal Pradesh, western Himalaya. *Current Science* **91(9)**: 1238-1242.
- Upadhye AS, Vartak VD and Kumbhojkar MS 1994. Ethno-medico-botanical studies in western Maharashtra, India. *Ethnobotany* **6(1-2)**: 25-31.
- Valvi SR, Deshmukh SR and Rathod VS 2011. Ethnobotanical survey of wild edible fruits in Kolhapur district. *International Journal of Applied Biology and Pharmaceutical Technology* **2(1)**: 194-197.