

Phytochemical characterization of Himalayan barberry (*Berberis lycium*) leaves

POONAM CHAUHAN* and ASHWANI TAPWAL

ICFRE – Himalayan Forest Research Institute, Shimla 171013 Himachal Pradesh, India

*Email for correspondence: chauhanp5396@gmail.com

© Society for Advancement of Human and Nature (SADHNA)

Received: 07.11.2025/Accepted: 12.12.2025

ABSTRACT

Berberis lycium Royle, commonly known as Himalayan barberry, is an important medicinal shrub traditionally used in the treatment of various ailments. The present study investigates the phytochemical composition of *B lycium* leaves using Gas Chromatography-Mass Spectrometry (GC-MS). Leaf samples collected from Solan district, Himachal Pradesh, were subjected to Soxhlet extraction using solvents of varying polarity: acetone, ethyl acetate and methanol. GC-MS profiling revealed a diverse phytochemical spectrum comprising polyketides, terpenoids, fatty acids, hydrocarbons, phenolics, phytosterols, vitamins and lipids. Ethyl acetate extract was rich in nonacosan-10-ol, phytol derivatives and tocopherols, while acetone extract predominantly contained γ -sitosterol, phenolics and hydrocarbons. Methanol extract exhibited high levels of polyketides, phytol, phytosterols and fatty acid derivatives. Several compounds such as phenols, alkaloids, flavonoids, lipids and lignans were documented, some of which are reported for the first time from *B lycium* leaves. The presence of these biologically active constituents highlights the therapeutic potential of the species and underscores its value for further pharmacological, biochemical and isolation studies.

Keywords: *Berberis lycium*; GC-MS; phytochemical analysis; leaf extract; bioactive compounds; phenolics; terpenoids; phytosterols

INTRODUCTION

Since ancient times, medicinal plants have been employed in traditional medical systems and the vast majority of people worldwide still rely on plants to treat a variety of diseases. Additionally, a range of plant-derived phytochemicals and their various parts are being studied for the treatment of many ailments, such as diabetes, cancer and many others. An estimated 70-80 per cent of people on the globe use medicinal plants to treat a variety of health issues (Ahmad et al 2024). *Berberis* is a large genus of evergreen and deciduous shrubs belonging to the family Berberidaceae, commonly known as barberry and is distributed across temperate and subtropical regions of the world. According to the Plant List this genus includes 628 species worldwide.

Berberis lycium Royle (Plate 1), one of the members of Berberidaceae family, is native to the Himalayan area, particularly Nepal, although it may also be found worldwide, in temperate and subtropical temperatures with the exception of Australia. John

Forbes Royle gave a description of *B lycium* in 1837 (Parra et al 2018). It is a highly valuable medicinal plant. In Hindi, this plant is known as Kashmal or Kasmal and in English as Indian barberry. It is a highly valuable medicinal plant that has been used for a very long time in conventional medicine (Ali et al 2015) and possesses broad spectrum antimicrobial activities against many pathogens (Parra et al 2018). It is a spiny shrub that grows to an average height of 2-4 metres, with lanceolate leaves arranged alternately on the stem. It bears yellow flowers in corymbose racemes and produces ovoid berries (Ahmed et al 2017). It is a source of many biologically active compounds. Phytochemicals of this plant include berbamine, berberine, jhelumine, palmatine, sindamine, punjabine, alkaloids, phenols, tannin, flavonoids, maleic acid, ascorbic acid, terpenoids, fats and resins (Anjum et al 2023). Singh and Gupta (2018) investigated antifungal activity of root, flower and fruit extract in different solvents (aqueous, ethanolic, Dimethyl formamide, propanol-2 and n-hexane) of *B lycium* against three fungal pathogens *Aspergillus niger*, *A fumigatus* and *A cuboida*. Their results revealed that ethanolic fruit

extract had significant antifungal activity than other extracts.

The objective of the present study was to assess the phytochemical composition of *B. lycium* by analysing extracts prepared using solvents of different polarities. The biologically active constituents in each extract were identified and characterized using Gas Chromatography-Mass Spectrometry (GC-MS). This study also provides a comprehensive profile of compounds that may be utilized in future research owing to their pharmacological properties.

MATERIAL and METHODS

Preparation of extract

The plant leaves were collected from Solan district, Himachal Pradesh, in the month of October. The samples were shade-dried and then ground into a fine powder. Soxhlet extraction was carried out using a thimble containing approximately 20 g of the powdered leaf material. Acetone, ethyl acetate and methanol were used as extraction solvents. Fig 1 illustrates the chromatograms of GC-MS of ethyl acetate leaf extract, acetone extract, methanol leaf extract of Himalayan barberry.

Phytochemical characterization

A GCMS system with an AOC-20i+s autosampler and a capillary column (HP-5MS, 30 m × 0.25 mm, 0.25 µm film thickness) was used to analyse the phytochemical components of the plant extract. Helium was used as the carrier gas at a constant flow rate of 1.0 ml/min. The injector temperature was set at 280°C and the column oven was initially maintained at 70°C for five minutes. A 1 µl sample was injected in splitless mode. After that, the oven's temperature was set to rise at a rate of 5°C per minute until 310°C, where it was maintained for ten minutes. The ion source and interface temperatures were kept at 230°C and 280°C respectively. With a scan range of m/z 50-700, the mass spectrometer was run in electron ionisation (EI) mode at 70 eV. The solvent cut time was set at 4.5 minutes and the whole run length was roughly 63 minutes. Spectral similarity indexes were used to compare the mass spectra of the compounds with the NIST libraries.

RESULTS

Phytochemical compounds identified and their molecular formula, weight, retention time, area

percentage are listed in the tables separately for each solvent.

A wide range of bioactive substances from several phytochemical groups were identified by GC-MS analysis of ethyl acetate leaf extract (Table 1).

Thirty four compounds, including polyketides, terpenoids, hydrocarbons, phenols, fatty acids, lipids and phytosterols, were found. Nonacosan-10-ol (30.05%), a lipid, was the most prevalent component and it was followed by 2-hexadecen-1-ol, 3,7,11,15-tetramethyl- (phytol acetate; 8.13%), dl- α -tocopherol (vitamin E, 7.66%) and quinic acid (7.10%). Phytol (5.22%), tricosyl acetate (4.46%), γ -sitosterol (4.01%) and pentacosane (4.06%) were other phytochemicals that add to the pharmacological relevance of the plant. Additionally, some compounds including 1,2-epoxyoctadecane (1.68%), n-hexadecanoic acid (1.65%) and neophytadiene (2.63%) were also found, suggesting the presence of terpenoid and fatty acid derivatives with possible bioactivity. Overall, the extract had a high concentration of hydrocarbons and lipid-based compounds; its aromatic and antioxidant profile was enhanced by other compounds such as o-cymene, γ -terpinene, thymol and 2,4-di-tert-butylphenol. The retention periods varied from 7.59 to 60.59 minutes, indicating a broad range of compound polarity. The phytochemical richness of *B. lycium* is shown by this chemical diversity, highlighting its potential as a valuable source of natural antioxidants, antimicrobials.

A comprehensive chemical profile comprising 23 bioactive chemicals from several chemical classes, such as alkanes, hydrocarbons, fatty acid derivatives, lipids, terpenes and phenolic compounds, was identified by GC-MS analysis of acetone extract (Table 2).

The most prevalent components were nonacosan-10-ol (7.34%), γ -sitosterol (8.30%) and 2,4-di-tert-butylphenol (9.38%). Tetradecane (4.57%), neophytadiene (3.82%), heptadecane (3.80%), 7,9-di-tert-butyl-1-oxaspiro (4,5)deca-6,9-diene-2,8-dione (3.26%) and eicosane (2.94%) are other significant chemicals that contribute to the pharmacological potential of the plant. Dodecane, hexadecane, phytol, and vitamin E are also important chemicals found in the extract contribute to its bioactivity. The retention time, which varied from 14.14 to 55.50 minutes, showed that chemicals with different polarity and volatility were present. The plant's vast phytochemical diversity is highlighted by the prevalence of alkanes and



Plate 1. *Berberis lycium* a. Flowering stage, b. Leaves




Fig 1. Chromatogram of GC-MS of Himalayan barberry a. Ethyl acetate leaf extract b. Acetone leaf extract c. Methanol leaf extract

hydrocarbons as well as the presence of phenolic and lipidic compounds.

A variety of bioactive substances, such as polyketides, amino acid derivatives, phenolics, terpenes, fatty acids and lipids, were found in the methanol extract according to the GC-MS analysis (Table 3).

2H-pyran-2,6(3H)-dione, which belongs to the polyketide class, was the most prevalent compound found (retention time 7.706 min, area 16.62%). Phytol (9.37%), 9,12,15-octadecatrienoic acid, methyl ester (Z,Z,Z) (9.33%), α -linolenic acid (6.01%), n-hexadecanoic acid (4.84%) and γ -sitosterol (5.56%) were among the significant compounds found. Butylsyringone (2.98%), dl-

Table 1. Phytochemicals present in the ethyl acetate extract of Himalayan barberry

Compound	Category	Molecular formula	Molecular mass	Retention time	Area (%)
2H-Pyran-2,6(3H)-dione	Polyketides	C ₅ H ₄ O ₃	112	7.595	1.92
o-Cymene	Monoterpenoid	C ₁₀ H ₁₄	134	8.636	0.12
γ-Terpinene	Monoterpene	C ₁₀ H ₁₆	136	9.785	0.15
1-Dodecene	Hydrocarbon	C ₁₂ H ₂₄	168	14.290	0.19
Thymol	Monoterpenoids	C ₁₀ H ₁₄ O	150	17.513	0.19
Phenol, 2-methoxy-3-(2-propenyl)	Phenol	C ₁₀ H ₁₂ O ₂	164	19.285	0.52
2,4-Di-tert-butylphenol	Phenol	C ₁₄ H ₂₂ O	206	23.506	0.46
1-Pentadecene	Hydrocarbons	C ₁₅ H ₃₀	210	25.663	0.53
Quinic acid	Cyclitol	C ₇ H ₁₂ O ₆	192	26.513	7.10
1-Octadecene	Hydrocarbons	C ₁₈ H ₃₆	252	30.386	0.53
Neophytadiene	Terpenoid	C ₂₀ H ₃₈	278	31.336	2.63
n-Hexadecanoic	Fatty acid	C ₁₆ H ₃₂ O ₂	256	34.008	1.65
1-Heptadecene	Hydrocarbons	C ₁₇ H ₃₄	238	34.651	0.41
Phytol	Diterpene	C ₂₀ H ₄₀ O	296	36.941	5.22
9,12,15-Octadecatrienoic acid	Fatty acid	C ₁₈ H ₃₀ O ₂	278	37.512	1.26
1-Hexacosene	Hydrocarbons	C ₂₆ H ₅₂	364	38.546	0.16
2-Hexadecen-1-ol,3,7,11,15-tetramethyl-, acetate	Phytol acetate	C ₂₂ H ₄₂ O ₂	338	38.881	8.13
3,7,11,15-Tetramethylhexadec-2-en-1-yl acetate	Terpenoids	C ₂₂ H ₄₂ O ₂	338	41.938	0.16
Heptacosane	Hydrocarbons	C ₂₇ H ₅₆	380	47.034	1.58
Pentacosane	Hydrocarbons	C ₂₅ H ₅₂	352	49.973	4.06
Tetracosan-10-ol	Fatty acid derivative	C ₂₄ H ₅₀ O	354	50.051	0.66
Octacosanal	Lipid	C ₂₈ H ₅₆ O	408	51.964	0.66
Pentacosane	Alkane	C ₂₅ H ₅₂	352	52.725	1.94
Nonacosan-10-ol	Lipids	C ₂₉ H ₆₀ O	424	52.848	30.05
dl-α-Tocopherol	Vitamin E	C ₂₉ H ₅₀ O ₂	430	53.175	7.66
Phetyl stearate	Fatty acid	C ₃₈ H ₇₄ O ₂	562	53.439	0.77
Tricosyl acetate	Fatty acid ester	C ₂₅ H ₅₀ O ₂	382	54.265	4.46
Triacontanal	Fatty acid derivative	C ₃₀ H ₆₀ O	436	54.801	0.70
1-Heptacosanol	Fatty alcohol	C ₂₇ H ₅₆ O	396	55.795	0.71
1,2-Epoxyoctadecane	Fatty acid derivative	C ₁₈ H ₃₆ O	268	56.031	1.68
γ-Sitosterol	Phytosterol	C ₂₉ H ₅₀ O	414	56.168	4.01
Oxirane, hexadecyl	Fatty acid derivative	C ₁₈ H ₃₆ O	268		
Clionasterol acetate	Phytosterol	C ₃₁ H ₅₂ O ₂	456	58.301	1.05
Hexadecanoic acid, 3,7,11,15-tetramethyl-2-hexadecenyl ester, [R-[R*,R*-(E)]]	Lipid	C ₃₆ H ₇₀ O ₂	534	60.591	0.67

α-tocopherol (2.85%), hexadecanoic acid, 2-hydroxy-1-(hydroxymethyl) ethyl ester (2.13%) and methyl syringate (2.02%) were other significant components. A complex mixture of phytochemicals with possible biological significance was found in moderate levels, including 5-

hydroxymethylfurfural (2.24%), dl-corydine (2.66%) and 9,12,15-octadecatrienoic acid, 2,3-dihydroxypropyl ester (3.02%). Trace amounts of minor chemicals were found, including L-proline 1-methyl, methyl ester, neophytadiene, sesamin, phenol, and 4-ethenyl-2,6-dimethoxy.

Table 2. Phytochemicals present in the acetone extract of Himalayan barberry

Compound	Category	Molecular formula	Molecular mass	Retention time	Area (%)
Dodecane	Alkane	C ₁₂ H ₂₆	170	14.143	3.05
2,4-Dimethylundecane	Alkane	C ₁₃ H ₂₈	184	16.067	1.07
2,6,11	Alkane	C ₁₅ H ₃₂	212	16.447	2.85
Trimethyldodecane					
Hexadecane	Alkane	C ₁₆ H ₃₄	226	17.853	0.71
Tetradecane	Alkane	C ₁₄ H ₃₀	198	20.133	4.57
Heneicosane	Alkane	C ₂₁ H ₄₄	296	20.534	0.51
Eicosane	Alkane	C ₂₀ H ₄₂	282	22.479	2.94
2,4-Di-tert-butylphenol	Phenol	C ₁₄ H ₂₂ O	206	23.081	9.38
Heptadecane	Hydrocarbon	C ₁₇ H ₃₆	240	25.379	3.80
3-Methylheptadecane	Hydrocarbon	C ₁₈ H ₃₈	254	29.391	0.32
Neophytadiene	Sesquiterpenoid	C ₂₀ H ₃₈	278	30.836	3.82
Phthalic acid, tetradecyl	Fatty acid	C ₃₂ H ₅₂ O ₄	500	31.373	1.49
trans-dec-3-enyl ester	derivatives				
7,9-Di-tert-butyl-1-oxaspiro(4,5)deca-6,9-diene-2,8-dione	Flavonoid	C ₁₇ H ₂₄ O ₃	276	32.346	3.26
Phytol	Terpene	C ₂₀ H ₄₀ O	296	36.447	1.25
triacontane, 1-iodo	Alkyl halide	C ₃₀ H ₆₁ I	148	41.699	0.88
Hexadecanoic acid, (2,2-dimethyl-1,3-dioxolan-4-yl) methyl ester	Fatty acid ester	C ₂₂ H ₄₂ O ₄	370	42.947	0.93
Dotriacontane, 1-iodo	Alkyl halide	C ₃₂ H ₆₅ I	576	43.360	0.50
Hexatriacontane	Hydrocarbon	C ₃₆ H ₇₄	506	44.980	0.53
Tetrapentacontane	Hydrocarbon	C ₅₄ H ₁₁₀	758	46.525	1.20
Hexadecanoic acid, eicosyl ester	Fatty acid	C ₃₆ H ₇₂ O ₂	536	46.525	0.93
2,6,10,14-Tetramethyloctadecane	Hydrocarbon	C ₂₂ H ₄₆	310	49.456	1.12
Nonacosan-10-ol	Lipid	C ₂₉ H ₆₀ O	424	52.308	7.34
Vitamin E	Vitamin	C ₂₉ H ₅₀ O ₂	430	52.661	1.59
γ-Sitosterol	Lipid	C ₂₉ H ₅₀ O	414	55.503	8.30

DISCUSSION

GC-MS profiling of *B lycium* leaves revealed a significant degree of phytochemical variation in solvent extracts of ethyl acetate, acetone and methanol. Nonacosan-10-ol, phytol derivatives, tocopherols and phenolic antioxidants were predominated in the ethyl acetate extract. The acetone extract was rich in γ-sitosterol, phenolics and hydrocarbons. Phytol, phytosterols, fatty acids (derivatives of linolenic acid) and polyketides (2H-pyran-2,6-dione) were abundantly present in the methanol extract. On the whole, the chemical composition was affected by the polarity of the solvent and the metabolites that were found showed significant biological properties such as antifungal, antibacterial, anti-cancerous, anti-inflammatory,

insecticidal, antiviral, antimalarial and antioxidant properties (Table 4) which make this plant economically important.

The effectiveness of *B lycium* in treating a variety of diseases has drawn attention, which has led researchers to embark on a number of experiments that aim at its pharmacological effects. This plant has historically been used to treat many diseases such as diarrhoea, intestinal colic, jaundice, piles, internal wounds, ophthalmia, rheumatism, and diabetes (Anjum et al 2023). According to experimental studies performed, *B lycium* was reported to have antibacterial, antihyperlipidemic, antidiabetic, anti-inflammatory, anticancer, antiviral and anthelmintic properties (Ahmad et al 2024). These diverse biological properties

Table 3. Phytochemicals present in the methanol extract of Himalayan barberry

Compound	Category	Molecular formula	Molecular mass	Retention time	Area (%)
2H-Pyran-2,6(3H)-dione	Polyketides	C ₅ H ₄ O ₃	112	7.706	16.62
L-Proline 1-methyl-, methylester	Proline derivative	C ₇ H ₁₃ NO ₂	143	9.231	0.47
Methylpyrrole-2-carboxylate	Organic acids	C ₆ H ₇ NO ₂	124	11.858	0.64
Butanedioic acid, hydroxy-, dimethyl ester (Malic acid)	Dicarboxylic acid	C ₆ H ₁₀ O ₅	162	12.025	0.67
5-Hydroxymethylfurfural (HMF)	Furan	C ₆ H ₆ O ₃	126	15.577	2.24
L-Proline, 5-oxo-, methyl ester (Pyroglutamic acid)	Amino acid derivative	C ₆ H ₉ NO ₃	143	20.207	1.91
Phenol, 4-ethenyl-2,6-dimethoxy	Methoxyphenols	C ₁₀ H ₁₂ O ₃	180	24.888	1.35
Cyclopentanol, 1,2-dimethyl-3-(1-methylethenyl)-, [1R-(1.alpha.,2.alpha.,3.alpha.)]	Terpene	C ₁₀ H ₁₈ O	154	27.225	0.23
Benzoic acid, 4-hydroxy-3,5-dimethoxy-, hydrazide	Phenol	C ₉ H ₁₂ N ₂ O ₄	212	29.828	0.53
1-Acetyl-2,2,6,6-tetramethyl-4-acetyloxypiperidine	Alkaloid	C ₁₃ H ₂₃ NO ₃	241	30.384	0.51
Butylsyringone	Phenolic	C ₁₂ H ₁₆ O ₄	224	31.000	2.98
Benzoic acid, 4-hydroxy-3,5-dimethoxy-, methyl ester (Methyl syringate)	Phenolic	C ₁₀ H ₁₂ O ₅	212	31.057	2.02
Neophytadiene	Terpene	C ₂₀ H ₃₈	278	31.332	1.93
Hexahydrofarnesyl acetone, 6,10,14-Trimethyl-2-pentadecanone	Sesquiterpene ketone	C ₁₈ H ₃₆ O	268	31.448	0.32
7,10,13-Hexadecatrienoic acid, methyl ester	Fatty acid	C ₁₇ H ₂₈ O ₂	264	32.609	0.49
7-Hexadecenoic acid, methyl ester	Fatty acid	C ₁₇ H ₃₂ O ₂	268	33.110	0.72
n-Hexadecanoic acid	Palmitic acid	C ₁₆ H ₃₂ O ₂	256	34.036	4.84
9,12-Octadecadienoic acid (Z,Z)-, methyl ester	Methyl linoleate	C ₁₉ H ₃₄ O ₂	294	36.608	1.76
9,12,15-Octadecatrienoic acid, methyl ester, (Z,Z,Z)	Methyl linolenate	C ₁₉ H ₃₂ O ₂	292	36.730	9.33
Phytol	Diterpene	C ₂₀ H ₄₀ O	296	36.946	9.37
9,12,15-Octadecatrienoic acid, (Z,Z,Z)	α -linolenic acid	C ₁₈ H ₃₀ O ₂	278	37.537	6.01
2-Hexadecen-1-ol, 3,7,11,15-tetramethyl-, acetate, [R-[R*,R*-(E)]]	Phytol acetate	C ₂₂ H ₄₂ O ₂	338	41.935	0.17
Hexadecanoic acid, 2-hydroxy-1-(hydroxymethyl) ethyl ester	Lipid	C ₁₉ H ₃₈ O ₄	330	44.079	2.13
9,12-Octadecadienoic acid (Z,Z)-, 2,3-dihydroxypropyl ester	Linoleoyl glycerol	C ₂₁ H ₃₈ O ₄	354	46.900	0.62
9,12,15-Octadecatrienoic acid, 2,3-dihydroxypropyl ester, (Z,Z,Z)	Fatty acid	C ₂₁ H ₃₆ O ₄	352	47.027	3.02
α -Tocospiro B	Terpenoid	C ₂₉ H ₅₀ O ₄	462	49.183	1.66
dl-Corydine	Benzylisoquinoline (Alkaloid)	C ₂₀ H ₂₃ NO ₄	341	49.461	2.66
Nonacosan-10-ol	Lipid	C ₂₉ H ₆₀ O	424	52.825	0.55
dl- α -Tocopherol	Polyterpene	C ₂₉ H ₅₀ O ₂	430	53.172	2.85
Phytol stearate	Terpenoids	C ₃₈ H ₇₄ O ₂	562	53.442	0.25
Sesamin	Lignan	C ₂₀ H ₁₈ O ₆	354	53.827	0.70
γ -Sitosterol	Lipids	C ₂₉ H ₅₀ O	424	56.177	5.56

Table 4. Compounds present in Himalayan barberry reported to have biological activities

Compound name	Biological activity	Reference
γ -Sitosterol	Anti-inflammatory, anti-cancer, cytotoxic, antihyperlipidemic, antidiabetic; treats ulcers, bronchitis, diabetes, heart diseases	Naikwadi et al (2022), Sonia and Singh (2019)
Sesamin	Antioxidant, anti-inflammatory, anticancer	Zhang et al (2022)
Phytol stearate	Found in plant wax	Garcés et al (2023)
dl- α -Tocopherol	Antioxidant	Serbinova et al (1991)
Nonacosan-10-ol	Component of tubular wax aggregates; also found in cuticular waxes of Pinaceae	Matas et al (2003), Dommissse et al (2007)
dl-Corydine	Inhibits growth of tumour cells	Konda et al (1990)
α -Tocospiro B	Antioxidant	Chen et al (2007)
Hexadecanoic acid, 2-hydroxy-1-(hydroxymethyl) ethyl ester	Anti-inflammatory, haemolytic; pesticidal, antioxidant	Tyagi and Agarwal (2017), Mohamed et al (2024)
2-Hexadecen-1-ol, 3,7,11,15-tetramethyl- acetate, [R-[R*, R*-(E)]]	Antibacterial	Thejashree and Naika (2023)
9,12,15-Octadecatrienoic acid, (Z,Z,Z)-	Antimicrobial, anticancer, hepatoprotective, anti-inflammatory	Banu and Nagarajan (2013)
Phytol	Antioxidant, anti-inflammatory, antimicrobial	Ashoka and Shivanna (2022)
9,12,15-Octadecatrienoic acid, methyl ester, (Z,Z,Z)	Antioxidant, anti-inflammatory, antibacterial, antifungal, anticancer	Mohamed et al (2015)
n-Hexadecanoic acid	Antimicrobial, antioxidant, hypocholesterolemic, nematocidal, pesticidal	Vedhanayaki and Ramkumar (2019)
Hexahydrofarnesyl acetone, 6,10,14-Trimethyl-2-pentadecanone	Antibacterial, anti-inflammatory, anticancer	Avoseh et al (2021)
Neophytadiene	Analgesic, antioxidant, antimicrobial, anticancer, antimalarial, insecticidal, neuroprotective	Rajeswaran and Rajan (2025)
1-Acetyl-2,2,6,6-tetramethyl-4-acetyloxypiperidine	Antimicrobial	Li et al (2015)
5-Hydroxymethylfurfural	Antioxidant, antiproliferative, cytotoxic	Zhao et al (2013)
2H-Pyran-2,6(3H)-dione	Antiallergic	Garst et al (1988)
Hexadecane	Antibacterial	Rouis-Soussi et al (2014)
Tetradecane	Antibacterial, antifungal	Naeim et al (2020)
Heneicosane	Antimicrobial	Vanitha et al (2020)
Vitamin E	Antioxidant	Brigelius-Flohé (2006)
o-Cymene	Insecticidal, insect repellent	Feng et al (2021)
γ -Terpinene	Antioxidant	Guo et al (2021)
Thymol	Antibacterial, antifungal	Marchese et al (2016)
Quinic acid	Antioxidant, antidiabetic, anticancer, antimicrobial, antiviral, antinociceptive, analgesic	Benali et al (2024)
2-Hexadecen-1-ol,3,7,11,15-tetramethyl-, acetate	Antibacterial	Thejashree and Naika (2023)
Heptacosane	Antioxidant, antibacterial, antimalarial, antitumor, antidermatophytic	Kuiate et al (2006)

of this plant are attributed to the presence of numerous phytochemicals.

Many studies were reported on the phytochemical characterization of root, flower and fruit by HPLC and qualitative analysis but very few on the leaf; therefore, this study has significance of documenting phytochemical compounds of leaves through GC-MS analysis for the first time.

Studies conducted on the phytochemical analysis of root confirmed the presence of steroids, alkaloids, flavonoids, tannins, terpenoids, anthoquinonine, proteins and carbohydrates (Mughal et al 2022) and through HPLC analysis presence of phenolic compounds (chlorogenic acid, quercetin, berberine, mandelic acid, rutin and hydroxy benzoic acid) were also confirmed (Nazir et al 2021).

GC-MS analysis of root, flower and fruit (methanolic extracts) were done by Gupta and Singh (2018) and they revealed the presence of some biologically active compounds, such as 3-benzyl-1,2-dihydronaphthalene, N-(4-biphenyl) benzamide, n-phenylbenzanilide etc. Stigmasterol was reported to be present in the root extract of *B lycium* by Sharma et al (2019) and also present in the leaves, in the current study.

Alkaloids, phenols, saponins, tannins, flavonoids, carbohydrates, terpenoids, steroids and glycosides were all found in *B lycium* leaves according to qualitative phytochemical examination and HPLC results have shown the presence of quercetin, gallic acid, ferulic acid and caffeic acid (Landry et al 2021).

In the current study, phenols (butylsyringone, benzoic acid, 4-hydroxy-3,5-dimethoxy-, methyl ester, 2,4-di-tert-butylphenol), alkaloids (dl-corydine, 1-acetyl-2,2,6,6-tetramethyl-4-acetyloxypiperidine), flavonoid (7,9-di-tert-butyl-1-oxaspiro(4,5) deca-6,9-diene-2,8-dione), terpenoids, lipids, vitamin, steroids, lipids and lignan etc are documented.

This study offers the first GC-MS-based phytochemical profiling of *B lycium* leaves. A number of compounds found in this analysis are being reported for the first time from this species and their biological activities are still mostly unknown. These results highlight the need for additional research, especially with regard to the isolation, structural characterisation and pharmacological evaluation of these newly

identified constituents in order to better understand their therapeutic potential.

CONCLUSION

This study investigates the presence of diversity of bioactive compounds with many of them reported to have biological properties according to the literature surveyed. Future investigations need to focus on in-depth mechanistic assessments that describe the molecular mechanisms by which *B lycium* exerts its effects, emphasising specific objectives for therapeutic development.

To assess its effectiveness and safety in humans, especially for its antidiabetic, anticancer and antiviral properties, carefully planned clinical trials are also required. To ensure consistency in therapeutic efficacy and safety, standardised extracts and formulations must be prepared. Using advanced techniques, further isolation and characterisation of the distinct phytochemicals found in this plant will bring insight into the particular compounds that are responsible of its pharmacological actions. Choosing the optimum growing techniques and plant traits for medicinal application can be facilitated by investigating the genetic and environmental factors influencing the phytochemical content and therapeutic properties of *B lycium*.

Despite the plant's widespread traditional use, only little research has investigated the chemical profile of leaves, rendering phytochemical characterisation essential. Determining the bioactive compounds affirms its ethnobotanical significance and establishes a scientific foundation for its medicinal properties. Additionally, this kind of characterisation aids in the selection of efficient compounds for focused pharmacological research. Further study on compound isolation, mechanism-of-action and the development of innovative medicinal formulations will be rendered feasible through the results presented here.

ACKNOWLEDGEMENTS

The authors express gratitude to Central Instrumentation Laboratory, Bhatinda, Punjab for GC-MS analysis and the financial support provided by the Council of Scientific and Industrial Research, New Delhi, India, under the CSIR research fellowship scheme (Grant Number 09/1346(12845)/2022-EMR-I).

REFERENCES

- Ahmad B, Murshed A, Zaky MY, Khan I, ur Rahman F, Abdellattif MH, Liu G and Lu J 2024. *Berberis lycium*: a miracle medicinal plant with multifaceted health benefits. *Journal of Food Quality* **2024**: 3152456; doi: 10.1155/2024/3152456.
- Ahmed S, Shuaib M, Ali K, Ali S and Hussain F 2017. Evaluation of different parts of *Berberis lycium* and their biological activities: a review. *Pure and Applied Biology* **6(3)**: 897-907.
- Ali H, Uddin S and Jalal S 2015. Chemistry and biological activities of *Berberis lycium* Royle. *Journal of Biologically Active Products from Nature* **5(5)**: 295-312.
- Anjum N, Ridwan Q, Akhter F and Hanief M 2023. Phytochemistry and therapeutic potential of *Berberis lycium* Royle; an endangered species of Himalayan region. *Acta Ecologica Sinica* **43(4)**: 577-584.
- Ashoka GB and Shivanna MB 2022. In vitro antioxidant and antibacterial activities of *Jatropha heynei* leaf extract and GC-MS profiling. *Asian Journal of Biological and Life Sciences* **11(3)**: 724-730.
- Avoseh ON, Mtunzi FM, Ogunwande IA, Ascrizzi R, Guido F 2021. *Albizia lebbek* and *Albizia zygia* volatile oils exhibit anti-nociceptive and anti-inflammatory properties in pain models. *Journal of Ethnopharmacology* **268**: 113676. doi: 10.1016/j.jep.2020.113676.
- Banu HR and Nagarajan N 2013. GC-MS determination of bioactive components of *Wedelia chinensis* (Osbeck) Merrill. *Journal of Chemical and Pharmaceutical Research* **5(4)**: 279-285.
- Benali T, Bakrim S, Ghchime R, Benkhaira N, El Omari N, Balahbib A, Taha D, Zengin G, Hasan MM, Bibi S and Bouyahya A 2024. Pharmacological insights into the multifaceted biological properties of quinic acid. *Biotechnology and Genetic Engineering Reviews* **40(4)**: 3408-3437.
- Brigelius-Flohé R 2006. Bioactivity of vitamin E. *Nutrition Research Reviews* **19(2)**: 174-186.
- Chen C-R, Chao L-H, Liao Y-W, Chang C-I, Pan M-H 2007. Tocopherols and triterpenoids from *Sida acuta*. *Journal of the Chinese Chemical Society* **54(1)**: 41-45.
- Dommissie A, Wirtz J, Koch K, Barthlott W and Kolter T 2007. Synthesis of (S) nonacosan 10 ol, the major component of tubular plant wax crystals. *European Journal of Organic Chemistry* **2007(21)**: 3508-3511.
- Feng Y-X, Zhang X, Wang Y, Chen Z-Y, Lu X-X, Du Y-S and Du S-S 2021. The potential contribution of cymene isomers to insecticidal and repellent activities of the essential oil from *Alpinia zerumbet*. *International Biodeterioration and Biodegradation* **157**: 105138; doi: 10.1016/j.ibiod.2020.105138.
- Garces R, de Andres-Gil C, Venegas-Calero M, Martinez-Force E, Moreno-Perez AJ and Salas JJ 2023. Characterization of sunflower seed and oil wax ester composition by GC/MS, a final evaluation. *LWT – Food Science and Technology* **173**: 114365; doi: 10.1016/j.lwt.2022.114365.
- Garst ME, Gluchowski C and Kaplan LJ 1988. Method for controlling the growth of microorganisms. US Patent Number 4,725,620, Patent and Trademark Office Washington, DC, US.
- Guo Y, Baschieri A, Amorati R, and Valgimigli L 2021. Synergic antioxidant activity of γ -terpinene with phenols and polyphenols enabled by hydroperoxyl radicals. *Food Chemistry* **345**: 128468; doi: 10.1016/j.foodchem.2020.128468.
- Gupta M and Singh A 2018. Pharmacological Studies of root, fruit and flower of *Berberis lycium*. *Oriental Journal of Chemistry* **34(2)**: 1055; doi: 10.13005/ojc/340257.
- Konda Y, Imai Y, Hojo H, Endo T and Nozoe S 1990. Suppression of tumor cell growth and mitogen response by aporphine alkaloids, dicentrine, glaucine, corydine and apomorphine. *Journal of Pharmacobiodynamics* **13(7)**: 426-431.
- Kuiate J-R, Bessière JM, Zollo PHA and Kuate SP 2006. Chemical composition and antidermatophytic properties of volatile fractions of hexanic extract from leaves of *Cupressus lusitanica* Mill from Cameroon. *Journal of Ethnopharmacology* **103(2)**: 160-165.
- Landry KB, Azam S, Rehman S, Tariq S, Iqbal B, Abbas M, Lembe DM and Ijaz B 2021. Phytochemical analysis of *Berberis lycium* methanolic extract and its antiviral activity through the restoration of MAPK signaling pathway modulated by HCV NS5A. *Asian Pacific Journal of Tropical Biomedicine* **11(3)**: 132-140.
- Li C, Hou J, Huang Z, Zhao T, Xiao L, Gao G, Harnode C and Dong A 2015. Assessment of 2,2,6,6-tetramethyl-4-piperidinol-based amine N-halamine-labeled silica nanoparticles as potent antibiotics for deactivating bacteria. *Colloids and Surface B: Biointerfaces* **126**: 106-114.
- Marchese A, Orhan IE, Daglia M, Barbieri R, Di Lorenzo A, Nabavi SF, Gortzi O, Izadi M and Nabavi SM 2016. Antibacterial and antifungal activities of thymol: a brief review of the literature. *Food Chemistry* **210**: 402-414.
- Matas AJ, Sanz MJ and Heredia A 2003. Studies on the structure of the plant wax nonacosan-10-ol, the main component of epicuticular wax conifers. *International Journal of Biological Macromolecules* **33(1-3)**: 31-35.

- Mohamed AA, Ali SI, Darwesh OM, El-Hallouty SM and Sameeh MY 2015. Chemical compositions, potential cytotoxic and antimicrobial activities of *Nitraria retusa* methanolic extract sub-fractions. *International Journal of Toxicology and Pharmacology Research* **7(4)**: 204-212.
- Mohamed HR, Hamed MM, El-Wakil EA and Okasha H 2024. GC-MS analysis, anti-inflammatory and anti-proliferative properties of the aerial parts of three *Mesembryanthemum* spp. *Toxicology Reports* **13**: 101829; doi: 10.1016/j.toxrep.2024.101829.
- Mughal TA, Ali S, Hassan A, Kazmi SAR, Saleem MZ, Shakir HA, Nazer S, Farooq MA, Awan MZ, Khan MA, Andleeb S, Mumtaz S, Mumtaz S, Tahir HM and Gulzar N 2022. Phytochemical screening, antimicrobial activity, in vitro and in vivo antioxidant activity of *Berberis lycium* Royle root bark extract. *Brazilian Journal of Biology* **84**: e249742; doi: 10.1590/1519-6984.249742.
- Naeim H, El-Hawiet A, Rahman RAA, Hussein A, El Demellawy MA and Embaby AM 2020. Antibacterial activity of *Centaurea pumilio* L root and aerial part extracts against some multidrug resistant bacteria. *BMC Complementary Medicine and Therapies* **20(1)**: 79; doi: 10.1186/s12906-020-2876-y.
- Naikwadi PH, Phatangare ND and Mane DV 2022. Active anti-inflammatory potency of γ -sitosterol from *Woodfordia floribunda* Salisb. *Journal of Plant Science Research* **38(2)**: 1-9.
- Nazir N, Rahman A, Uddin F, Khalil AAK, Zahoor M, Nisar M, Ullah S, Ullah R, Ezzeldin E and Mostafa GAE 2021. Quantitative ethnomedicinal status and phytochemical analysis of *Berberis lycium* Royle. *Agronomy* **11(1)**: 130; doi: 10.3390/agronomy11010130.
- Parra SA, Gaur K, Ranawat LS and Rather MI 2018. An overview on various aspects of plant *Berberis lycium* Royale. *American Journal of Pharmacological Sciences* **6(1)**: 19-24.
- Rajeswaran S and Rajan DK 2025. Neophytadiene: biological activities and drug development prospects. *Phytomedicine* **143**: 156872; doi: 10.1016/j.phymed.2025.156872.
- Rouis-Soussi LS, El Ayeb-Zakhama A, Mahjoub A, Flamini G, Jannet HB and Harzallah-Skhiri F 2014. Chemical composition and antibacterial activity of essential oils from the Tunisian *Allium nigrum* L. *EXCLI Journal* **13**: 526-535.
- Serbinova E, Kagan V, Han D and Packer L 1991. Free radical recycling and intramembrane mobility in the antioxidant properties of alpha-tocopherol and alpha-tocotrienol. *Free Radical Biology and Medicine* **10(5)**: 263-275.
- Sharma P, Awasthi A and Gupta I 2019. Phytochemical investigation and evaluation of anti-oxidant potential of *Berberis lycium* roots from Himachal Pradesh. *Journal of Pharmacognosy and Phytochemistry* **8(3)**: 4740-4745.
- Singh A and Gupta M 2018. Phytochemical investigation of different solvent extracts of *Berberis lycium* fruits. *Rasayan Journal of Chemistry* **11(1)**: 228-231.
- Sonia and Singh S 2019. Phytoconstituents of *Ziziphura nummularia* (Burm f) Wight and Arn leaves extracts using GC-MS spectroscopy. *Research and Reviews: A Journal of Life Sciences* **9(1)**: 109-118.
- Thejashree AB and Naika R 2023. Identification of bioactive compounds in acetone leaf and stem-bark extracts of *Psychotria dalzellii* Hook f by GC-MS analysis and evaluation of in vitro anti-bacterial properties. *Asian Journal of Biological and Life sciences* **12(3)**: 499-509.
- Tyagi T and Agarwal M 2017. Ethnomedicine analysis of bioactive constituents in ethanolic leaf extract of *Pistia stratiotes* L and *Eichhornia crassipes* (Mart) Solms by GC-MS. *Advances in Bioresearch* **8(5)**: 204-211.
- Vanitha V, Vijayakumar S, Nilavukkarasi M, Punitha VN, Vidhya E and Praseetha PK 2020. Heneicosane – a novel microbicidal bioactive alkane identified from *Plumbago zeylanica* L. *Industrial Crops and Products* **154**: 112748; doi: 10.1016/j.indcrop.2020.112748.
- Vedhanayaki TC and Ramkumar B 2019. Analysis of bioactive compounds in ethanolic extract of *Commelina maculata* leaves using GC-MS technique. *Journal of Pharmacognosy and Phytochemistry* **8(4)**: 867-870.
- Zhang Y, Liu F, Lin Y, Li L, Chen M and Ni L 2022. A comprehensive review on distribution, pharmacological properties and mechanisms of action of sesamin. *Journal of Chemistry* **2022**: 4236525; doi: 10.1155/2022/4236525.
- Zhao L, Chen J, Su J, Li L, Hu S, Li B, Zhang X, Xu Z and Chen T 2013. In vitro antioxidant and antiproliferative activities of 5-hydroxymethylfurfural. *Journal of Agricultural and Food Chemistry* **61(44)**: 10604-10611.