

EFFECT OF POTASSIUM NITRATE ON YIELD AND QUALITY OF CHILLI

Babanjeet*, Kulbir Singh, Dilpreet Talwar and Salesk Kumar Jindal

Department of Vegetable Science, Punjab Agricultural University, Ludhiana-141004, Punjab

ABSTRACT

The present investigation was conducted during 2019-20 and 2020-21 to study the effect of potassium nitrate on yield and quality of chilli. The experiment was conducted in randomized complete block design and replicated thrice during both the years. The experiment comprised of twelve treatment combinations of four nitrogen doses (0, 75, 100 and 125 kg N/ha) and three potassium levels (0, 30 kg K/ha and 30 kg K/ha along with foliar spray of KNO_3 @ 2%). Two equal splits of nitrogen i.e., one at the time of transplanting and remaining at first picking were applied in the plot where 75 kg/ ha N was incorporated. However, for 100 and 125 kg N/ha, application was made in three equal splits i.e., first at the time of transplanting, second and third at first and third picking of chilli. Foliar application of KNO_3 was made thrice i.e., at the time of flowering and after first and third picking of chilli. Potassium nitrate (KNO_3) @ 2% along with 100 kg N/ha and 30 kg K/ha proved to be significantly higher in growth parameters such as canopy width and number of branches; yield parameters such as fruit length, fruit diameter, green fruit yield (q/ha), red fruit yield (q/ha); quality parameters such as dry matter, colouring matter, capsaicin content in green and red chilli, and oleoresin content as compared to recommended dose of fertilizer (RDF) comprising 75 kg N/ha and 30 kg K/ha. Higher green and red ripe fruit yield of 8-18 % and 20-22 %, respectively in chilli was recorded due to an increase in above mentioned parameters as compared to RDF during both years. Similarly, higher dry matter (16-20%) and capsaicin content (6-9%) were recorded under same treatments. So, 100 kg nitrogen per hectare in three splits and 30 kg potassium per hectare along with three foliar sprays of KNO_3 @ 2% in chilli proved to be suitable for production of chilli under Punjab conditions.

Keywords: Capsaicin Content, Chilli, Foliar Application, Fruit Yield, Potassium Nitrate

Indian cuisine is renowned and celebrated throughout the world for its spicy treat to the tongue. An unforgettable experience has been created by the spices which impart flavor and aroma to the food. Among the commonly utilized spices in Indian food, chilli is an inevitable ingredient (Saxena, 2016). Chilli (*Capsicum annuum* var. *annuum* L.) is an important spice and condiment crop grown almost everywhere in the world. It thrives well in tropical and sub-tropical region (Dhaliwal, 2015). The chromosome number of chilli is $2n=24$ and it belongs to the family Solanaceae (Naz *et al.*, 2006). Chilli is also known for its medicinal effects and therapeutic uses like blood circulation, cholesterol and blood pressure maintenance, hence beneficial in heart disorders (Saleh *et al.*, 2018). It is also used as a condiment in various food items like curry sauces, soups and chutney to add flavour and pungency; moreover, it is widely used in the preparation of various kinds of pickles. Green chilli is eaten raw in salads as well as cooked. Chilli powder is made up of 54% pericarp, 40% seed and 6% stalk (Prasad *et al.*, 2016). Chilli contains numerous biochemical substances like alkaloid (capsaicin), vitamins, organic acids, minerals and flavonoids which are thought to

be anti-carcinogenic by having antioxidant properties (Hasler and Clare, 1998). Chilli covers an area of 7.43 lac and 8.78 thousand hectares with production of 19.14 lac and 17.63 thousand tonnes and with productivity of 2576 kg/ha and 20.09 q/ha in India (Anonymous, 2021a) and Punjab, respectively (Anonymous 2021b).

Nutrients play an essential role in plant growth and development. Various nutrients have different biochemical and physiological roles in plant system. Nitrogen is a chief constituent of various important substances which occur in plants like protein, chlorophyll and nucleic acids (Singh and Kumar, 1999). Nitrogen application to chilli peppers exhibited an important role for rise in plant growth parameters, color and nutritive value of leaves and yield (Tumbare and Nikram, 2004). The crucial role of nitrogen for being an integral part of all amino acids in proteins and lipids, the structural complexes of cells and chloroplast made it the most important macronutrient for better plant formation and proper growth (Uddin *et al.*, 2003).

Potassium is also considered to be an essential nutrient supplying 6% of plant dry weight (Hartz *et al.*, 1998). Potassium is not a component of any organic molecule but, it plays a chief role in various biochemical and physiological processes i.e., osmoregulation process, regulation of stomata opening and closing, water use efficiency, translocation of sugars and

*Corresponding author : babanjeet-vs@pau.edu
Date of receipt: 30.03.2022, Date of acceptance: 31.12.2022

formation of carbohydrates which effects crop quality, size of fruit and stimulates root growth (Chauhan, 2012). It is also called a quality nutrient because it directly enhances quality attributes of plants. Potassium nitrate is a water soluble potassic fertilizer containing 44% K₂O and 13% N suitable for foliar fertilization. The foliar spray of potassium nitrate in chilli considerably increases the growth and yield parameters by influencing various physiological and biochemical activities. In vegetable crop production, fertilization has a significant role to play, not only the nutrient but its source and method of application is also important. In future, foliar nutrient sprays will be of great utility because it reduces the total amount of fertilizer applied, increases the availability of trace elements and high efficiency of fertilizer application is also achieved (Haytova *et al.*, 2013). Lester *et al.* (2006) pointed out that insufficient information regarding subsistence farming and conservative attitude like application of an inadequate dose of potassium in soil to fulfill the crop requirement for increased yields were the reasons for less availability of nutrients.

Chilli is a long duration crop, hence it requires an adequate supply of manures and fertilizers. To provide adequate nutrition for proper plant growth, the physiological stage of development must be studied. The application of different combinations of N & K fertilizers at fruit maturity and development may improve red colour, oleoresin content, pungency, growth, yield and quality (Ananthi *et al.*, 2004). Therefore, there is a fundamental need to develop a nutrient management strategy for chilli that will not only improve nutrient use efficiency but also increase quality and yield of the produce. So, keeping in the view importance of potassium nitrate in enhancing quality and yield of various vegetables, the proposed investigation was conducted to determine the optimum concentration of potassium nitrate for better growth, quality and yield of chilli.

MATERIALS AND METHODS

The investigation was conducted at the Research farm of Vegetable Science and Biochemistry Laboratory, PAU, Ludhiana during 2019-20 and 2020-21. The experiment was conducted in randomized complete block design and replicated thrice. The nursery of chilli crop (CH 27) was sown in mid-November of 2019 and 2020 which was transplanted in mid-February 2020 and 2021 for two years. The crop was irrigated by furrow irrigation system. Rest of cultural practices were followed as per standard agronomic practices (Anonymous, 2021b). The experiment comprised of twelve treatment combinations of four nitrogen (0, 75, 100 and 125 kg N/ha) and three potassium (0, 30, 30 kg K/ha along with foliar spray of KNO₃ @ 2%). Two equal splits of nitrogen i.e., one at the time of transplanting and remaining at first picking were applied in the plot where 75 kg/ha N

was incorporated. However, for 100 and 125 kg N/ha application was made in three equal splits i.e., first at the time of transplanting, second and third at first and third picking of chilli. Foliar application of KNO₃ was made thrice i.e., at the time of flowering, first and third picking of chilli. Growth related parameters i.e., Canopy width (cm), number of branches and yield related parameters i.e., Fruit length (cm), fruit diameter (cm), green & red ripe fruit yield (g/plant), fruit yield (q/ha) as well as quality parameters like dry matter, capsaicin content were estimated by method recommended by Bajaj and Kaur (1979) and oleoresin were estimated as per method suggested by Tandon *et al.* (1964).

RESULTS AND DISCUSSION

Growth parameters

Canopy width (cm)

Canopy width is an important parameter depicting the health of the plant. The data pertaining to the effect of different treatments on canopy width of chilli in all directions at final harvest is presented in Table 1. Data showed that maximum canopy width after final harvest was observed with treatment T₁₂ (125 kg/ha of N (Three split doses) + 30 kg/ha of K + 3 foliar spray of KNO₃ @ 2%) which was significantly higher than all other treatments. On the other hand, minimum canopy width was witnessed under control (T₁). The findings are in consonance with the observations of Khan *et al.* (2011) that higher dose of nitrogen resulted in higher canopy width as compared to other lower doses. Similarly, Stewart *et al.* (2003); Wahocho *et al.* (2015) observed that incorporation of inorganic sources of nitrogen and potassium enhances canopy width in chilli. Here, higher nitrogen and potassium in plants mean higher rates of accumulation of photosynthates, which accelerates canopy width of the plant.

Number of branches per plant

Ibrahim (2003) reported that number of branches per plant affect fruit yield of plant both directly and indirectly. It is an important trait which regulates fruit bearing ability of plant. Sometimes an association was found between number of branches, number of leaves, photosynthetic activity and LAI which promotes higher bearing ability of plant. The data pertaining to the effect of different treatments of nitrogen and potassium on number of branches per plant is presented in Table 1. Maximum number of branches per plant (7.33) were recorded with treatment T₁₀ (125 kg/ha of N + 0 kg/ha of K) during 2019-20 and 2020-2021, which was significantly higher from all of other treatments. Treatments T₃, T₇, T₁₁ and T₁₂ was recorded to be statistically at par with each other in 2019-2020. Whereas, in pooled data number

Table 1. Effect of different treatments on canopy width (cm) and number of branches per plant

S. No.	Treatment	Canopy width (cm)			Number of branches per plant		
		2019-20	2020-21	Mean	2019-20	2020-21	Mean
T ₁	0 kg/ ha of N + 0 kg/ ha of K	47.12	48.07	47.6	5.78	5.69	5.73
T ₂	0 kg/ha of N + 30 kg/ ha of K	54.57	56.35	55.46	6.40	6.62	6.51
T ₃	0 kg/ ha of N + 30 kg/ ha of K + 3 foliar sprays of KNO ₃ @ 2%	58.70	59.48	59.09	6.67	6.38	6.52
T ₄	75 kg of N (Two split doses) + 0 kg/ha of K	60.90	60.41	60.65	6.13	6.36	6.25
T ₅	75 kg of N (Two split doses) + 30 kg/ha of K	62.90	61.79	62.34	6.04	6.23	6.14
T ₆	75 kg of N (Two split doses) + 30 kg/ha of K + 3 foliar sprays of KNO ₃ @ 2%	70.02	71.85	70.94	5.96	6.00	5.98
T ₇	100 kg/ha of N (Three split doses) + 0 kg/ha of K	66.52	69.8	68.16	6.98	6.73	6.85
T ₈	100 kg/ha of N (Three split doses) + 30 kg/ha of K	73.70	78.28	75.99	6.44	6.68	6.56
T ₉	100 kg/ha of N (Three split doses) + 30 kg/ha of K + 3 foliar sprays of KNO ₃ @ 2%	80.12	79.22	79.67	6.36	6.63	6.49
T ₁₀	125 kg/ha of N (Three split doses) + 0 kg/ha of K	70.29	74.46	72.38	7.33	7.00	7.17
T ₁₁	125 kg/ha of N (Three split doses) + 30 kg/ha of K	83.84	85.91	84.87	6.84	6.73	6.79
T ₁₂	125 kg/ha of N (Three split doses) + 30 kg/ha of K + 3 foliar sprays of KNO ₃ @ 2%	92.29	91.71	92.00	7.16	6.92	7.04
CD (P = 0.05)		6.76	8.91	5.50	0.88	0.23	0.47

of branches per plant with application of recommended dose of fertilizer (T₅) was recorded about 6.14, which was significantly higher from control T₁ (5.73). These findings are in accordance with the findings of Mohanty *et al.* (2001); Gare *et al.* (2001) showed that increasing the amount of nitrogen and potassium up to certain level helped to increase the number of branches per plant which in turn increased the fruit yield. A comparative relationship occurs between nitrogen application and accumulation of biomass in plants. So higher concentration of nitrogen results in higher content of biomass which further increases the number of branches.

Yield parameters

Fruit length

Fruit length is an important parameter in chilli. Fruit length is positively associated with fruit yield. Minimum fruit length was obtained with no fertilizer application (T₁). Whereas maximum fruit length was observed with treatment T₉ (100 kg/ ha of N (Three split doses) + 30 kg/ ha of K + 3 foliar spray of KNO₃@ 2%) which was significantly higher from rest of the treatments during both seasons. In pooled data, the fruit length attained under treatment T₉ was 12% more than that obtained with recommended dose (T₅) and 9.6% more than that obtained with no fertilizer application. But different treatments failed to show any significant effect on fruit length of chilli during first season. Data related to

this finding is shown in Table 2. Similar findings were observed in experiment of Ananthi *et al.* (2004); Lodhi *et al.* (2019) in which surge in fruit length was observed with the rise in nitrogen and potassium application. Here, fruit length increased with the accumulation of biomass in plant which in turn is related with the concentration of nitrogen and potassium present in plant system.

Fruit diameter

Fruit diameter is positively related to the fruit yield. It is important for determining the actual size of the fruit. It also portrays the health of the fruit and consumption of secondary metabolites by plant. Data pertaining to fruit diameter is shown in the Table 2. Treatment T₉ (100 kg/ ha of N (Three split doses) + 30 kg/ ha of K + 3 foliar spray of KNO₃@ 2%) resulted in maximum fruit diameter (1.33 cm) which was significantly higher from rest of the treatments in pooled data. However, all the treatments failed to show any noteworthy effect on fruit diameter of chilli during both seasons. Fruit diameter obtained with control (T₁) was recorded lower than recommended dose. The results are in conformity with the findings of Golcz *et al.* (2012) which showed that there was decrease in fruit diameter with the increase in the level of potassium. Similarly, Hartz *et al.* (2001) found that soil drenched with higher dose of potassium did not affect fruit diameter significantly. The purpose behind this finding is that a high level of potassium results in high production of organic acids that leads to osmotic imbalance and adverse effects on nutrition

Table 2. Effect of different treatments on Fruit length (cm), Fruit diameter (cm), Green fruit yield (g/plant), Green fruit yield (q/ha), Red fruit yield (g/plant), Red fruit yield (q/ha) of chilli.

S. No.	Treatment	Fruit length (cm)			Fruit diameter (cm)			Green fruit yield (g/plant)			Green fruit yield (q/ha)			Red fruit yield (g/plant)			Red fruit yield (q/ha)		
		2019-20	2020-21	Pooled data	2019-20	2020-21	Pooled data	2019-20	2020-21	Pooled data	2019-20	2020-21	Pooled data	2019-20	2020-21	Pooled data	2019-20	2020-21	Pooled data
T ₁	0 kg/ha of N + 0 kg/ha of K	6.43	6.29	6.36	1.28	1.24	1.26	1351.80	1337.70	1344.80	206.00	201.40	203.70	914.30	912.60	913.40	188.10	176.50	182.30
T ₂	0 kg/ha of N + 30 kg/ha of K	6.46	6.31	6.39	1.27	1.23	1.25	1355.30	1356.70	1356.00	206.50	213.60	210.10	968.60	966.70	967.70	199.30	187.60	193.40
T ₃	0 kg/ha of N + 30 kg/ha of K + 3 foliar sprays of KNO ₃ @ 2%	6.52	6.32	6.42	1.28	1.28	1.28	1484.90	1493.30	1489.10	226.30	229.90	228.10	1013.30	1016.50	1014.90	208.50	204.00	206.20
T ₄	75 kg of N (Two split doses) + 0 kg/ha of K	6.47	6.38	6.43	1.27	1.28	1.28	1373.20	1406.70	1389.90	209.30	215.60	212.40	1000.80	991.20	996.00	205.90	193.60	199.80
T ₅	75 kg of N (Two split doses) + 30 kg/ha of K	6.54	6.58	6.56	1.27	1.30	1.29	1516.50	1527.90	1522.20	231.10	252.70	241.90	1023.60	1037.80	1030.70	210.60	218.70	214.60
T ₆	75 kg of N (Two split doses) + 30 kg/ha of K + 3 foliar spray of KNO ₃ @ 2%	6.84	6.81	6.83	1.31	1.29	1.30	1679.30	1670.00	1674.70	255.90	265.50	260.70	1156.60	1181.30	1169.00	237.90	252.20	245.10
T ₇	100 kg/ha of N (Three split doses) + 0 kg/ha of K	6.62	6.43	6.52	1.29	1.28	1.29	1581.50	1573.90	1577.70	241.00	235.90	238.50	1093.30	1086.50	1089.90	224.90	228.70	226.80
T ₈	100 kg/ha of N (Three split doses) + 30 kg/ha of K	6.63	6.54	6.58	1.29	1.29	1.29	1596.30	1596.60	1596.40	243.30	245.90	244.60	1098.10	1095.50	1096.80	225.90	242.90	234.40
T ₉	100 kg/ha of N (Three split doses) + 30 kg/ha of K + 3 foliar sprays of KNO ₃ @ 2%	6.94	7.45	7.19	1.31	1.35	1.33	1796.70	1787.10	1791.90	273.80	273.40	273.60	1233.30	1242.30	1237.80	253.80	263.80	258.80
T ₁₀	125 kg/ha of N (Three split doses) + 0 kg/ha of K	6.61	6.39	6.50	1.29	1.32	1.31	1525.80	1491.70	1508.70	232.50	232.70	232.60	1040.00	1081.30	1060.60	213.90	222.50	218.20
T ₁₁	125 kg/ha of N (Three split doses) + 30 kg/ha of K	6.68	6.68	6.68	1.28	1.30	1.29	1634.00	1610.00	1622.00	249.00	246.30	247.70	1124.50	1146.00	1135.20	231.60	248.20	239.80
T ₁₂	125 kg/ha of N (Three split doses) + 30 kg/ha of K + 3 foliar sprays of KNO ₃ @ 2%	6.84	7.00	6.92	1.29	1.31	1.30	1712.50	1725.60	1719.00	260.90	251.00	256.00	1196.70	1232.40	1214.50	246.20	253.60	249.90
CD (P = 0.05)		NS	0.08	0.20	NS	NS	NS	155.20	111.40	99.40	23.60	17.40	15.20	113.90	143.10	95.40	17.40	21.80	14.20

translocation in plant system which resulted in non-significant effect on fruit diameter.

Green fruit yield

The yield is also an important criterion for comparing the efficiency of different treatments. Yield potential is the deciding factor of adaptability of any crop plant. The data regarding the effect of different treatments on green fruit yield (g/plant) is presented in Table 2. Maximum yield (g/plant) was attained with T₉ (100 kg/ ha of N + 30 kg/ ha of K + 3 foliar spray of KNO₃@ 2%) which was significantly higher than all other treatments during both the seasons. In pooled data of two seasons, the maximum green fruit yield per plant (1791.87 g) was observed in treatment T₉ (100 kg/ ha of N + 30 kg/ ha of K + 3 foliar spray of KNO₃@ 2%) which was significantly higher by 17.71% and 33.25% from recommended dose of fertilizers and control, respectively. The results are in conformity with the findings of Solangi *et al.* (2013) that significant green fruit yield can only be achieved with definite dose of potassium and nitrogen. Similarly, Prasad *et al.* (2016) also found that various doses of nitrogen and potassium affect green fruit yield differently.

Red fruit yield

Red fruit yield (g/plant) is the measure of total quantity of red fruits produced. During both seasons and in pooled data the maximum yield recorded with T₉ (100 kg/ ha of N + 30 kg/ ha of K + 3 foliar spray of KNO₃@ 2%) was significantly higher than rest of the treatments. During 2019-20, treatment T₅, T₆, T₇, T₈, T₉, T₁₁, T₁₂ and during 2020-21 treatment T₆, T₈, T₁₁ and T₁₂ were found to be statistically at par with each other. The results are in conformity with the findings of Dubey *et al.* (2017) that increasing dose of nitrogen and potassium resulted in increasing total red yield of chilli. Similar results were found by Hiremath *et al.* (2006); Ortas (2013) in pepper and tomato, respectively.

Quality parameters

Dry matter

Dry matter is the sum total of all constituents of plant excluding water. It is the result of spatial and temporal integration of all plant activities. In case of crop canopies studies, it is the most applicable parameter. The data related to the effect of different nitrogen and potassium fertilizer treatments on dry matter of chilli is presented in Table 3. Perusal of data showed that highest dry matter content (28.80% and 28.50%) during both seasons was recorded in T₉ (100 kg/ ha of N + 30 kg/ ha of K + 3 foliar spray of KNO₃@ 2%). However, in both seasons dry matter values attained with different treatments (T₆, T₇, T₈, T₉, T₁₁ and T₁₂) were found to be statistically similar with each other. Same result was observed in pooled

data of two seasons. These results are in conformity with Pervez *et al.* (2004) which showed a promoting effect of nitrogen was reported on accumulation of dry matter in chilli crop. Accumulation of dry matter depends upon the photosynthetic rate of plant which further depends upon the chlorophyll content.

Colouring matter (ASTA)

Capsanthin and capsorubin are the two main carotenoids pigments which are primarily responsible for colour development in chilli. These pigments impart orange to red colour in the fruit (Kendrick, 2012). In Table 3, pooled data showed that with T₉ (100 kg/ ha of N + 30 kg/ ha of K + 3 foliar spray of KNO₃@ 2%) there is about 6.50 % and 9.37% more colouring matter as compared to recommended dose of fertilizers (T₅) and control (T₁). In both seasons 2019-20 and 2020-21 as well as in pooled data of two seasons maximum colouring matter (240.1) was attained with the application of T₉ which was significantly better than rest of the treatments. However, during 2019-20 data has revealed that treatments T₆, T₉, T₁₁ and T₁₂ were found to be statistically similar with each other. Same results have been observed during 2020-21 as well as in pooled data of both seasons. The results were in accordance with the findings of Bidari *et al.* (2004) where increased concentration of colouring matter was observed with increased nitrogen and potassium application. However, it was noted that potassium dose had much higher effect on the content of colouring matter than nitrogen. The reason is that potassium regulates the availability of substrates which are responsible for synthesis of capsanthin and capsorubin pigments.

Capsaicin content in green chilli

In Table 3 data has showed that the maximum capsaicin content in green chilli was attained with the application of T₉ (100 kg/ ha of N + 30 kg/ ha of K + 3 foliar spray of KNO₃@ 2%) during 2019-20 (0.5390 %) and 2020-21 (0.5300 %) as well as in pooled data of two seasons (0.5345 %) which was significantly better than rest of all treatments. In pooled data with treatment T₉ (100 kg/ ha of N + 30 kg/ ha of K + 3 foliar spray of KNO₃@ 2%), 16.37 % and 24.68 % more capsaicin content in green chilli was recorded than recommended dose of fertilizers (T₅) and control (T₁) respectively; similar results were found by Data and Jana, 2009.

Capsaicin content in red chilli

Pungency of chilli is due to capsaicinoids and active compound of chilli called capsaicin. Capsaicin is mainly restricted to placenta and pericarp of chilli fruit which has various physiological functions. In Table 3 pooled data showed that with T₉ (100 kg/ ha of N + 30 kg/ ha of K + 3 foliar spray of KNO₃@ 2%) 7.48 % and

Table 3. Effect of different treatments on Dry matter, colouring matter (ASTA units), Capsaicin content in green chilli (%), Capsaicin content in red chilli (%), Oleoresin content (%) of chilli.

S. No.	Treatment	Dry matter (%)			Colouring matter (ASTA units)			Capsaicin content in green chilli (%)			Capsaicin content in red chilli (%)			Oleoresin content (%)		
		2019-20	2020-21	Pooled data	2019-20	2020-21	Pooled data	2019-20	2020-21	Pooled data	2019-20	2020-21	Pooled data	2019-20	2020-21	Pooled data
T ₁	0 kg/ha of N + 0 kg/ha of K	20.70	20.57	20.64	220.8	218.3	219.6	0.430	0.427	0.429	0.6833	0.6827	0.6830	9.13	9.15	9.14
T ₂	0 kg/ha of N + 30 kg/ha of K	21.47	21.67	21.57	222.4	222.6	222.5	0.438	0.434	0.436	0.6983	0.6977	0.6980	9.30	9.37	9.33
T ₃	0 kg/ha of N + 30 kg/ha of K + 3 foliar sprays of KNO ₃ @ 2%	23.67	23.87	23.77	224.8	223.6	224.2	0.456	0.453	0.454	0.7190	0.7183	0.7187	11.01	11.07	11.04
T ₄	75 kg of N (Two split doses) + 0 kg/ha of K	22.02	22.27	22.15	223.9	222.8	223.4	0.448	0.446	0.447	0.7040	0.7093	0.7067	10.63	10.43	10.53
T ₅	75 kg of N (Two split doses) + 30 kg/ha of K	24.63	24.37	24.50	225.9	224.8	225.4	0.461	0.458	0.459	0.7290	0.7287	0.7288	11.48	11.49	11.49
T ₆	75 kg of N (Two split doses) + 30 kg/ha of K + 3 foliar spray of KNO ₃ @ 2%	27.70	27.53	27.62	238.7	234.3	236.5	0.485	0.482	0.483	0.7650	0.7640	0.7645	12.65	12.55	12.60
T ₇	100 kg/ha of N (Three split doses) + 0 kg/ha of K	26.20	25.97	26.09	228.6	226.8	227.7	0.469	0.467	0.468	0.7450	0.7447	0.7448	11.64	11.48	11.56
T ₈	100 kg/ha of N (Three split doses) + 30 kg/ha of K	26.47	26.30	26.39	230.6	229.6	230.1	0.476	0.471	0.474	0.7520	0.7510	0.7515	12.06	12.18	12.12
T ₉	100 kg/ha of N (Three split doses) + 30 kg/ha of K + 3 foliar spray of KNO ₃ @ 2%	28.80	28.50	28.65	240.7	239.4	240.1	0.539	0.530	0.535	0.7830	0.7833	0.7833	13.50	13.39	13.45
T ₁₀	125 kg/ha of N (Three split doses) + 0 kg/ha of K	25.03	24.47	24.75	226.9	225.6	226.2	0.466	0.462	0.464	0.7343	0.7337	0.7340	11.59	11.54	11.57
T ₁₁	125 kg/ha of N (Three split doses) + 30 kg/ha of K	27.07	26.83	26.95	233.6	232.3	232.9	0.479	0.476	0.478	0.7580	0.7573	0.7577	11.89	11.70	11.80
T ₁₂	125 kg/ha of N (Three split doses) + 30 kg/ha of K + 3 foliar spray of KNO ₃ @ 2%	28.33	28.20	28.27	237.9	236.8	237.3	0.510	0.495	0.503	0.7687	0.7680	0.7683	12.74	12.42	12.58
CD (P = 0.05)		2.74	2.68	2.20	9.4	10.9	8.4	0.009	0.008	0.0078	0.0030	0.0031	0.0028	1.31	0.94	0.76

14.69 % more capsaicin content was obtained in red chilli as compared to recommended dose of fertilizers (T_5) and control (T_1). However, maximum capsaicin content (0.7833 %) was attained with the application of T_9 (100 kg/ ha of N + 30 kg/ ha of K + 3 foliar spray of KNO_3 @ 2%) which was significantly better than rest of the treatments. Veerana *et al.* (2000) showed that increase in potassium content significantly increase the capsaicin content in chilli fruit. The reason behind the finding is nitrogen in plant system helps to form various amino acids. Branched amino acids like valine and leucine plays the primary role in the synthesis of capsaicin (Sarma *et al.*, 2004).

Oleoresin

Deep red coloured semi viscous liquid obtained from chilli is called as oleoresin. It is used in various food industries as a flavouring agent, colouring agent and preservative (Harp and Barrows, 2015). The data presented in Table 3 revealed that T_9 (100 kg/ ha of N + 30 kg/ ha of K + 3 foliar spray of KNO_3 @ 2%) during both seasons 2019-2020 and 2020-21 as well as pooled data resulted in maximum oleoresin content (13.45%) which was significantly higher from recommended dose of fertilizers (T_5) 11.49% and control (T_1) 9.14%. This result is comparable with the findings of Prabhavathi *et al.* (2008) that higher level of potassium nitrate application brought about maximum oleoresin content in chilli fruit. Ananthi *et al.* (2004) in an experiment observed that nitrogen and potassium also increased oleoresin content in chilli upto certain level despite showing the increment in other growth factors.

From whole study, it can be concluded that the growth parameters such as canopy width and number of branches; yield parameters such as fruit length, fruit diameter, green fruit yield(q/ha), red fruit yield (q/ha); quality parameters such as dry matter, colouring matter, capsaicin content in green and red chilli, oleoresin content were higher due to interactive effect of KNO_3 as compared to RDF alone. To conclude, application of potassium nitrate (KNO_3) @ 2% along with 100 kg/ ha of N (Three split doses) and 30 kg/ ha of K showed overall better performance in growth, quality and yield of chilli than top dressed fertilizers alone.

Authors' contribution

Conceptualization of research work and designing of experiments (BJ, KS, DT, SJ); Execution of field/lab experiments and data collection (BJ, KS, SJ); Analysis of data and interpretation (BJ, KS, DT); Preparation of manuscript (BJ, KS, DT, SJ).

LITERATURE CITED

Ambare T P, Gonge V S, Deshmukh D T, Warade A D and Hameed B 2005. Effect of nitrogen on growth and yield of

Okra (*Abelmoschus esculentus*) varieties. *J Maharashtra Agric Univ* **30**: 217-18.

Aminifard M H, Aroiee H, Fatemi H and Ameri A 2010. Performance of eggplant and sweet pepper in intercropping system under different rates of nitrogen. *Hortic Environ Biotechnol* **51**: 367-72.

Ananthi S, Veeraragavathatham D and Srinivasan K 2004. Comparative efficiency of muriate of potash and sulphate of potash on yield attributes, yield and economics of chilli (*Capsicum annum* L.). *Indian J Hort* **52**: 158-63.

Anonymous 2021a. *Horticultural statistics at a glance*. Retrived from National Horticultural Board, pp. 202.

Anonymous 2021b. *Package of practices for vegetable crops*. Punjab Agricultural University, Ludhiana, pp. 46-49.

Bajaj K L and Kaur G 1979. Colorimetric determination of capsaicin in capsicum fruits with the Folin-Ciocalteu reagent. *Mikrochim Acta* **71**: 81-86.

Bidari B I, Martur M D and Math K K 2004. Influence of soil properties on yield and quality of chillies (*Capsicum annum* L.) and partitioning of nutrients in fruit components. *Asian J Soil Sci* **1**: 49-51.

Chauhan D V S 2012. *Vegetable production*. Kalyani Publications, New Delhi, 85p.

Data S and Jana J C 2009. Effect of seasons and fertility levels on yield and quality of chilli genotypes in terai-agroclimatic region. *Indian J Agric Sci* **79**: 917-21.

Dhaliwal M S 2015. *Handbook of Vegetable Crops*. Kalyani Publications, India, 274p

Dubey A K, Rajput D S, Singh P, Verma Y K, Kumar A and Chandraker S K 2017. Effect of NPK on plant growth, yield and quality of capsicum cv Swarna under shade net conditions. *Int J Curr Microbiol Appl Sci* **6**: 1085-91.

Gare B N, More S M, Jadhav M G and Burlia V 2001. Effect of spacing and fertilizer on yield of rainfed chilli in sub-montane zone of Maharashtra. *J Maharashtra Agric Univ* **25**: 270-71.

Golcz A, kujawski P and Markiewicz B 2012. Yielding of red pepper under the influence of varied potassium fertilization. *Sci Hort* **11**: 3-15.

Harp B P and Barrows J N 2015. US regulation of colour additives in foods. In: scotter, M.J *Colour Additives for Foods and Beverages* (eds.). Woodhead publishing, Oxford, pp 25-40

Hartz T K, Miyao E M and Valencia J G 1998. DRIS evaluation of the nutritional status of processing tomato. *Hortic Sci* **33**: 830-32.

Hartz T K, Miyao G, Mullen, Cohn M D, Valencia J and Brittan K L 2001. Potassium requirements for maximum yield and fruit quality of tomato. *J Am Soc Hort Sci* **124**: 199-204.

Hasler M and Clare P H 1998. Functional foods and their role in disease prevention and health promotion. *J Food Technol* **52**: 63-147.

- Haytova D, Kostova D and Mechandjiev D 2013. Effect of type and method of fertilization on marrows (*Cucurbita pepo* L.) yield and fruit quality. *J Exp Agric Int* **4**: 376-83.
- Hiremath S M, Basavraj N and Dharmathi R R 2006. Response to location Spacing and fertilizers levels on yield and yield attributes of paprika. *J Agric Sci* **19**: 362-65.
- Ibrahim R 2003. Correlation studies and percent contributions of different growth and yield characters to total fruit yield of tomato at Samaru, Nigeria. *J Hortic Sci* **26**: 463 (Abstr).
- Kendrick A 2012. Natural food and beverage colorings. In: Baines D and Seal R *Natural Food Additives, Ingredients and flavouring* (eds.). Woodhead publishing, Cambridge, pp. 25-40.
- Khan Z, Tiyaqi S A, Mahmood I and Rizvi R 2011. Effects of N fertilization organic matter and biofertilisers on the growth and yield of chilli in relation to management of plant-parasitic nematodes. *J Bot* **36**: 73-81.
- Lester G E, Jifon J L and Makus D J 2006. Supplemental foliar potassium applications with or without a surfactant can enhanced vegetable quality. *Hortic Sci* **41**: 741-44.
- Lodhi Y, Chakravorty S, Prasad BVG and Sangeeta 2019. Influence of nutrients and mulching on fruiting and fruit characteristics of bell pepper. *J Pharm Innov* **8**: 791-94.
- Mohanty B K, Hossain M M and Nanda S K 2001. Response of chilli cv. Utkal Rashmi to nitrogen, phosphorus and potash. *Orissa J Hortic* **29**: 51-53.
- Naz S, Anjum MA and Ahmad I 2006. Growth of chilli (*Capsicum annuum* L.) F1 hybrid sky line-2 in response to different ages of transplants. *J Res Sci* **17**: 91-95.
- Ortas I 2013. Influence of nitrogen and potassium fertilizer rates on pepper (*Capsicum annum* L.) and tomato (*Lycopersicon esculentum*) yield and nutrient uptake under field conditions. *Prog Hort* **8**: 1048-55.
- Pervez M A, Ayub C M, Bashart A, Nave A V and Nasir M 2004. Effect of nitrogen levels and spacing on growth and yield of chilli (*Capsicum annum* L.). *Int J Agric Biol* **6**: 504-06.
- Prabhavathi K, Bidari B I, Shashidhara G B and Mathad J C 2008. Influence of sources and levels of potassium on quality attributes and nutrient composition of red chillies (*Capsicum annum* L.). *J Agric Sci* **21**: 379-81.
- Prasad S R, Singh K and Chandra H 2016. Effect of phosphorus and organic matter on dry matter, yield and nutrients uptake in chilli (*Capsicum annum* L.). *Ann Agri Bio Res* **18**: 315-21.
- Reddy P V, Jyothi K U, Bharathi S and Suryakumari S 2013. Effect of nitrogen and potassium sources on yield and yield attributes of chilli (*Capsicum annum* L.). *J Spices Aromat Crops* **24**: 137-40.
- Sajid M, Ishtiaq M, Amin N, Arif, M and Rehman Z 2001. Effect of nitrogen and phosphorus on the growth and yield of red chillies (*Capsicum annum* L.). *Sarhad J Agric* **17**: 549-51.
- Saleh B K, Omer A, Teweldemedhin B 2018. Medicinal uses and health benefits of chilli pepper (*Capsicum spp.*) *J Food Process Technol* **6**: 325-28.
- Sarma U J, Baruah J P, Suhrawardy J, Narzry B D and Chakravarty M 2004. Effect of various NPK levels on yield and capsaicin content in direct seeded chilli (*Capsicum annum* L.). *Indian J Hill Agric* **17**: 15-18.
- Shivkumar M S, Nagaraja B V and Kuligod V B 2010. Comparative analysis of different levels of nutrient management in dry chilli production- a case study. *Indian J Fertilizers* **6**: 12-16.
- Singh P and Kumar A 1999. Genetic variability, heritability and genetic advances in chilli. *Indian J Agric Sci* **77**: 459-61.
- Solangi A H, Baloch P A, Abro B A and Siddiqui A 2013. Growth and yield characteristics of chilli (*Capsicum annum* L.) as affected by N in presence and absence of P and K. *Pak J Sci Ind Res B Biol Sci* **56**: 70-75.
- Stewart D W, Costa C, Dwyer L M, Smith D L and Hamilton R I 2003. Canopy structure, light interception and photosynthesis in maize. *J Agron* **95**: 1465-74.
- Tandon G L, Dravid S V and Siddappa G S 1964. Oleoresin of capsicum and some technological and chemical aspects. *J Food Sci* **29**: 1-5.
- Tumbare A D and Nikram A D 2004. Effect of planting and fertigation on growth and yield of green chilli. *Indian J Agri Sci* **74**: 242-45.
- Uddin K, Khalequzzaman K M, Rahman M, Siddique N A and Ali O 2003. Yield and yield components of winter chilli as affected by different levels of nitrogen and boron. *Pak J Biol Sci* **6**: 605-09.
- Veerana H K, Abdul K, Farooqui A A and Sujith G M 2000. Effect of potassium fertigation on growth and yield of chilli (*Capsicum annum* L.) under moisture stress conditions. *J Spices Aromat Crops* **29**: 117-21.
- Wahocho N A, Sheikh Z A, Jogi Q, Talpur Q and Leghari S J 2015. Growth and productivity of chilli under various nitrogen levels. *Curr Sci Int* **28**: 1321-26.